

THE COS/UVES SURVEY OF THE MAGELLANIC STREAM: II. EVIDENCE FOR A COMPLEX ENRICHMENT HISTORY OF THE STREAM FROM THE FAIRALL 9 SIGHTLINE

PHILIPP RICHTER^{1,2}, ANDREW J. FOX³, BART P. WAKKER⁴, NICOLAS LEHNER⁵, J. CHRISTOPHER HOWK⁵, JOSS
BLAND-HAWTHORN⁶, NADYA BEN BEKHTI⁷, CORA FECHNER¹

¹Institut für Physik und Astronomie, Universität Potsdam, Haus 28, Karl-Liebknecht-Str. 24/25, 14476 Golm (Potsdam), Germany

²Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany

³Space Telescope Science Institute, Baltimore, MD 21218, USA

⁴Department of Astronomy, University of Wisconsin-Madison, 475 N. Charter Street, Madison, WI 53706, USA

⁵Department of Physics, University of Notre Dame, 225 Nieuwland Science Hall, Notre Dame, IN 46556, USA

⁶Institute of Astronomy, School of Physics, University of Sydney, NSW 2006, Australia and

⁷Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

Draft version from March 26, 2013

ABSTRACT

We present a multi-wavelength study of the Magellanic Stream (MS), a massive gaseous structure in the Local Group that is believed to represent material stripped from the Magellanic Clouds. We use ultraviolet, optical and radio data obtained with *HST*/COS, VLT/UVES, *FUSE* and GASS to study metal abundances and physical conditions in the Stream toward the quasar Fairall 9. Line absorption in the MS is detected from a large number of metal ions and from molecular hydrogen (H_2). The UVES data indicates a complex velocity-component structure with seven absorption components related to the MS. From the analysis of unsaturated Si II absorption, in combination with a detailed photoionization model, we obtain a surprisingly high α abundance in the Stream toward Fairall 9 of $[\text{S}/\text{H}] = -0.30 \pm 0.04$ (0.50 solar). This value is 5 times higher than what is found along other MS sightlines based on similar COS/UVES data sets. In contrast, the measured nitrogen abundance is found to be substantially lower ($[\text{N}/\text{H}] = -1.15 \pm 0.06$), implying a very low $[\alpha/\text{N}]$ ratio of -0.85 dex. The elements Si, Fe, Ni, Ti, Al are depleted into dust grains with depletion values in the range $\delta \approx -0.5$ to -1.6 dex. The H_2 absorption in the Stream is accompanied by very narrow optical Na I/Ca II absorption. Our analysis of these species suggests the presence of a cold ($T \sim 100$ K) and dense ($n_{\text{H}} \approx 5 \text{ cm}^{-3}$) gas phase in the body of the Stream, possibly arranged in a widespread population of pc-scale clumps or filaments. From the analysis of C II* absorption we infer a local electron density of $n_{\text{e}} \leq 0.05 \text{ cm}^{-3}$ and a mean C⁺ cooling rate of $7.3 \times 10^{-27} \text{ erg s}^{-1}$. The substantial differences in the chemical composition of the Magellanic Stream toward Fairall 9 compared to other lines of sight, together with the presence of dust and molecular gas, point toward a complex enrichment history of the Magellanic Stream. We favour a scenario, in which the gas was locally enriched with α elements by massive stars in a star-forming region in the SMC or LMC and then was expelled and/or stripped from the stellar disk before the delayed nitrogen enrichment from intermediate-mass stars could set in.

Subject headings: ISM: abundances – Galaxy: halo – Galaxy: evolution – Magellanic Clouds – quasars: absorption lines

1. INTRODUCTION

The distribution of neutral and ionized gas in the circumgalactic environment of galaxies is known to be an important indicator of the past and present evolution of galaxies. Both the infall of metal-poor gas from intergalactic space and from satellite galaxies and the outflow of metal-rich gaseous material through galactic winds represent key phenomena that determine the spatial distribution and the physical state of the circumgalactic gas around massive galaxies.

From observations and theoretical studies, it is known

that galaxy interactions between gas-rich galaxies can transport large amounts of neutral and ionized gas into the circumgalactic environment of galaxies. In the local Universe, the most massive of these extended tidal gas features can be observed in the 21 cm line of neutral hydrogen (HI). The most prominent nearby example of a tidal gas stream produced by the interaction of galaxies is the Magellanic Stream (MS), a massive ($\sim 10^8 - 10^9 M_{\odot}$) stream of neutral and ionized gas in the outer halo of the Milky Way at a distance of $\sim 50 - 60$ kpc (e.g., Wamier & Wrixon 1972; Gardiner & Noguchi 1996; Weiner & Williams 1996; Putman et al. 2003; Brüns et al. 2005; Fox et al. 2005, 2010; Koerwer 2009; Besla et al. 2007, 2010, 2012). The Magellanic Stream spans over 200° on the sky (e.g., Nidever et al. 2010) and has a (mean) metallicity that is lower than that of the Milky Way, but comparable with the metallicity found in the SMC and

¹ Based on observations obtained with the NASA/ESA Hubble Space Telescope, which is operated by the Space Telescope Science Institute (STScI) for the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5D26555, and on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere, Chile.

LMC ($0.1 - 0.5$ solar; Lu et al. 1994; Gibson et al. 2000; Sembach et al. 2001; Fox et al. 2010, 2013). The MS also contains dust grains and diffuse molecular hydrogen (H_2 ; Sembach et al. 2001; Richter et al. 2001a). A number of theoretical studies, including tidal models and ram-pressure stripping models, have been carried out to describe the Stream’s motion in the extended halo of the Milky Way and pinpoint its origin in one of the two Magellanic Clouds (Gardiner & Noguchi 1996; Mastrogiro et al. 2005; Connors et al. 2006; Besla et al. 2010; Diaz & Bekki 2011).

The origin and fate of the Magellanic Stream is closely related to the trajectories of LMC and SMC (e.g., Connors et al. 2004, 2006; Besla et al. 2007), and any realistic model of the MS thus needs to consider the dynamical and physical state of the Milky Way/Magellanic Clouds system as a whole (see also Bland-Hawthorn et al. 2007). While early tidal models have assumed that the Magellanic Stream is a product from the tidal interaction between LMC and SMC as they periodically orbit the Milky Way (e.g., Gardiner & Noguchi 1996), more recent proper motion measurements of the Magellanic Clouds (MCs) (Kallivayalil et al. 2006a, 2006b) indicate that the MCs may be on their first passage around the Milky Way. Some subsequent tidal models (Besla et al. 2010; Diaz & Bekki 2011) thus favour a first-infall scenario for the Magellanic Stream. Moreover, while many models (e.g., Connors et al. 2006) place the origin of the Stream’s gaseous material in the SMC, other, more recent studies trace back at least part of the Stream’s gaseous material in the LMC (e.g., Nidever et al. 2008). The latter study also highlights the role of energetic blowouts from star-forming regions in the LMC for the formation of the Stream. Clearly, further theoretical studies and observations are required to pinpoint the origin of the MS based on different (independent) methods.

In the first paper in our series analyzing the chemical and physical conditions in the Magellanic Stream (Fox et al. 2013; hereafter Paper I), we have investigated MS absorption in the UV and optical along the lines of sight toward RBS 144, NGC 7714 PHL 2525, and HE 0056–3622. In this paper we analyze the MS using UV and optical absorption-line spectra of the Seyfert 1 galaxy Fairall 9 ($z_{\text{em}} = 0.047$). Located at $l = 295.1$ and $b = -57.8$ the Fairall 9 sightline lies only 14.3° on the sky from the SMC. This sightline is the best-studied in absorption of all MS directions (Songaila 1981; York et al. 1982; Lu, Savage & Sembach 1994; Gibson et al. 2000; Richter et al. 2001a; Sembach et al. 2003), largely because the Fairall 9 is bright in both the optical and the UV and the Stream’s H I column in this direction is large ($\log N(\text{H I}) \approx 20$; see Gibson et al. 2000). The high column of neutral gas ensures that a wide range of low-ionization UV metal lines are detectable in the Stream, and even molecular hydrogen was observed in the MS toward Fairall 9 data from the *Far Ultraviolet Spectroscopic Explorer* (*FUSE*; Richter et al. 2001a; Wakker 2006). Using a spectrum of Fairall 9 obtained with the Goddard High Resolution Spectrograph (GHRS) onboard the *Hubble Space Telescope* (HST) together with Parkes 21 cm H I data Gibson et al. (2000) derived a metallicity of the Stream toward Fairall 9 of $[\text{S/H}] = -0.55 \pm 0.06^{+0.17}_{-0.21}$ (~ 0.3 solar), which represented the most accurate metal-

licity determination of the Stream from UV absorption line data at that time. This metallicity is consistent with either an SMC or LMC origin of the gas. A difficulty for constraining the origin the MS in one or the other Magellanic Cloud arises from the fact that the gas in the Stream was stripped from its parent galaxy $\sim 1 - 2$ Gyr ago (e.g., Gardiner & Noguchi 1996; Connors et al. 2006; Nidever et al. 2008), but has not experienced any further metal enrichment since then, while the parent galaxy underwent further chemical evolution. The MS does not contain any massive stars (e.g., Mathewson et al. 1979), in contrast to the Magellanic Bridge (Irwin, Kunkel & Demers 1985). This aspect needs to be taken into account when comparing metal abundances in the Stream with present-day LMC and SMC abundances.

To increase the accuracy of the metallicity determination of the Stream toward Fairall 9 and to obtain more detailed information on the chemical composition of the gas and dust in the Stream, more accurate spectral data are desirable. Because the Magellanic Stream is a massive gas cloud with complex internal kinematics (e.g., Nidever et al. 2008), data with high spectral resolution and a high signal-to-noise (S/N) ratio are required to fully resolve the Stream’s velocity-component structure and to detect weak absorption features from the various metal ions that have their transitions in the ultraviolet (UV) and in the optical. As part of our ongoing project to study the properties of the Magellanic Stream in absorption along multiple lines of sight (see also Paper I) we obtained high-resolution optical data of Fairall 9 from the Ultraviolet and Visible Echelle Spectrograph (UVES) installed on the Very Large Telescope (VLT) and medium-resolution UV data from the Cosmic Origins Spectrograph (COS) onboard the *HST*, both data sets providing absorption spectra with excellent S/N ratios. The combination of these data sets, as described in this study, therefore provides a particular promising strategy to study in great detail the chemical and physical conditions in the Magellanic Stream in this direction.

This paper is organized as follows: in Sect. 2 we describe the observations and the data reduction. The column density measurements and the profile modeling are explained in Sect. 3. In Sect. 4 we derive chemical and physical properties of the gas in the MS. We discuss our results in Sect. 5. Finally, a summary our study is given in Sect. 6.

2. OBSERVATIONS AND SPECTRAL ANALYSIS

2.1. VLT/UVES observations

Fairall 9 was observed with the VLT/UVES spectrograph (Dekker et al. 2010) in 2010 under ESO program ID 085.C-0172(A) (PI: A. Fox). The observations were taken in Service Mode using Dichroic 1 in the $390+580$ setting, a $0.6''$ slit, and no rebinning. The observations were carried out under good seeing conditions ($< 0.8''$). The raw data were reduced with the standard UVES pipeline, using calibration frames taken close in time to the corresponding science frames. The reduction steps involve subtraction of the bias level, inter-order background, sky background, night sky emission lines, and cosmic ray hits. The frames were then flat-fielded, optimally extracted and merged. The wavelength scale was corrected for atmospheric dispersion and heliocentric ve-

locity and then placed into the local standard of rest (LSR) velocity frame. Multiple exposures on the same target were registered onto a common wavelength grid and then added. The final spectra have a very high spectral resolution of $R \approx 70\,000$ corresponding to a FWHM of 4.3 km s^{-1} . They cover the wavelength range between 3300 and 6800 Å. The S/N ratio per resolution element is 40 at 3500 Å (Ti II), 65 at 4000 Å (Ca II), and 83 at 6000 Å (Na I). The UVES data thus provide much higher sensitivity and substantially higher spectral resolution than previous optical measurements of Fairall 9 (Songaila 1981).

2.2. HST/COS observations

Fairall 9 was observed with the *HST*/COS spectrograph (Green et al. 2012) in 2012 under *HST* program ID 12604 (PI: A. Fox). A four-orbit visit provided a total of 5378 s of exposure time with the G130M/1291 wavelength setting, and 6144 s with the G160M/1589 setting. With each grating, all four FP-POS positions were used to dither the position of the spectrum on the detector to reduce the fixed-pattern noise. The raw data were processed and combined with the CALCOS pipeline (v2.17.3). For the coaddition of the individual exposures we used interstellar absorption lines as wavelength reference. The final, co-added spectra then were transformed into the LSR velocity frame. The COS data cover the UV wavelength range between 1131 and 1767 Å. The spectra have a resolution of $R \approx 16\,000$ (FWHM $\approx 19\text{ km s}^{-1}$) and a pixel size of $\sim 7\text{ km s}^{-1}$. The S/N ratio per resolution element is 37 at 1200 Å and 26 at 1550 Å.

In order to minimize geocoronal emission, which contaminates the absorption lines of O I $\lambda 1302$, Si II $\lambda 1304$ and H I Ly α in the velocity range $-200\text{ km s}^{-1} \lesssim v_{\text{LSR}} \lesssim +200\text{ km s}^{-1}$ during orbital daytime, we re-reduced the data with a night-only extraction. For this, data were extracted from those time intervals when the Sun's altitude was less than 20 degrees.

2.3. FUSE observations

As part of our study we also re-analyze archival *FUSE* spectra of Fairall 9. These spectra were obtained in 2000 under *FUSE* program ID P101 (PI: K.R. Sembach) with a total exposure time of 34827 s. They show strong molecular hydrogen absorption arising in the MS, as presented by Richter et al. (2001a). The *FUSE* spectra have a resolution of $\sim 20\text{ km s}^{-1}$ (FWHM), and cover the wavelength range 912 – 1180 Å. The raw data were reduced using the CALFUSE pipeline v3.2.1. The individual exposures were carefully coadded using interstellar lines as wavelength reference. Unfortunately, the S/N in the *FUSE* data is very low, which severely hampers the analysis of absorption lines (see also Richter et al. 2001a; Wakker 2006). From a detailed inspection of the individual spectra in the different segments we conclude that only the data from the LiF 1A segment can be properly coadded without introducing large systematic errors in the flux distribution in the spectrum. In the coadded LiF 1A data, that are used by us to re-analyze the H₂ absorption in the MS, we measure a S/N of ~ 6 per resolution element at 1020 Å. Only for the wavelength range 1050 – 1082 Å does the S/N ratio rise up to a




FIG. 1.— H I column-density map of the Magellanic Stream in the direction of Fairall 9, based on 21cm data from GASS. The map shows the distribution of neutral gas in the LSR velocity range between +100 and +250 km s^{-1} . The position of Fairall 9 is indicated with the white ellipse, whose shape reflects the beam size of the radio observations from GASS.

maximum of ~ 15 per resolution element due to the increased background flux from the broad, redshifted Ly β emission from Fairall 9.

2.4. GASS 21cm observations

The H I 21cm data for the Fairall sightline were taken from the Galactic-All Sky Survey (GASS, McClure-Griffiths et al. 2009, Kalberla et al. 2010). The survey was observed with the 64-m radio telescope at Parkes. The data cubes have an angular resolution of 15/6, leading to an RMS of 57 mK per spectral channel ($\Delta v = 0.8\text{ km s}^{-1}$). This value translates to an H I column density detection limit of $N(\text{H I})_{\text{lim}} = 4.1 \times 10^{18}\text{ cm}^{-2}$, assuming a Gaussian-like emission line with a width of 20 km s^{-1} FWHM. Fig. 1 shows a H I column density map of the local environment of the Magellanic Stream centered on Fairall 9.

2.5. Spectral analysis methods

Our strategy for the analysis of the optical and UV absorption-line data of Fairall 9 combines different techniques to optimally account for the different spectral resolutions and S/N in the data. The reduced and coadded spectra from UVES, COS, and *FUSE* first were continuum-normalized using low-order polynomials that were fitted locally to the spectral regions of interest.

For the high-resolution VLT/UVES data we have used Voigt-profile fitting to decompose the MS absorption pattern in the optical lines of Ca II, Na I, and Ti II into individual absorption components (Voigt components) and to derive column densities (N) and Doppler parameters (b values) for the individual components. For the fitting, we made use of the FITLYMAN package implemented in the ESO-MIDAS analysis software (Fontana & Ballester 1995). Laboratory wavelengths and oscillator strengths have been taken from the compilation of Morton (2003). Note that with our fitting technique we are able to measure b values smaller than the instrumental resolution as we *simultaneously* fit the line doublets of Ca II and Na I, so that relative strengths of the lines are taken into account for the determination of both b and N . From




FIG. 2.— Optical absorption profiles of Ca II, Na I, and Ti II from VLT/UVES data of Fairall 9 are shown. Absorption from the Magellanic Stream is seen at LSR velocities between +130 and +240 km s^{-1} . The solid red line displays the best possible Voigt profile fit to the data. The individual velocity components are indicated by the tick marks. In the upper panel, the H I 21cm emission spectrum toward Fairall 9 from the GASS data is plotted for comparison.

this fitting procedure we obtain a component model for Fairall 9 sightline, in which the LSR velocity centroids of the individual absorption components and the b values for the low ions are defined.

In addition to Voigt-profile fitting, we have used the apparent optical depth method (AOD method; Savage & Sembach 1991) to derive total gas column densities for the (unsaturated) optical absorption profiles of Ca II, Na I, and Ti II. The AOD analysis was made using the custom-written MIDAS code *span* that allows us to measure equivalent widths and AOD column densities (and their errors) in absorption spectra from a direct pixel integration.

For the medium-resolution *HST*/COS data we have used profile *modeling* and the AOD method to derive column densities and column density limits for the various different low, intermediate, and high ions that have detectable transitions in the COS wavelength range. The ion transitions in the COS data considered in this study include C II $\lambda\lambda 1334.5, 1335.7$, C IV $\lambda\lambda 1548.2, 1550.8$, N I $\lambda\lambda 1199.6, 1200.2, 1200.7$, N V $\lambda\lambda 1238.8, 1242.8$, O I $\lambda 1302.2$, Al II $\lambda 1670.8$, Si II $\lambda\lambda 1190.4, 1193.3, 1260.4, 1304.4, 1526.7$, Si III




FIG. 3.— Velocity profiles of various low ions in the COS data of Fairall 9 are shown. The FUV data for Si II $\lambda 1020.7$ are from *FUSE*. The velocity components related to gas in the Magellanic Stream, as identified in the optical Ca II data, are indicated with the tick marks. The red solid line indicates the best-fitting absorption model for the UV absorption, as described in Sect. 3.2. The single, saturated lines of C II $\lambda 1334.5$ and O I $\lambda 1302.2$ are not considered in our model.

$\lambda 1206.5$, Si IV $\lambda\lambda 1393.8, 1402.8$, P II $\lambda 1152.8$, S II $\lambda\lambda 1250.6, 1253.8, 1259.5$, Fe II $\lambda\lambda 1143.2, 1144.9, 1608.5$, and Ni II $\lambda 1370.1$. For the profile modeling of neutral and singly-ionized species that trace predominantly neutral gas in the MS we have used as input the component model defined by the optical Ca II absorption (see above). In this model, the LSR velocities and b values of the seven absorption components are constrained by the Ca II fitting results, while the column density for each component is the main free parameter that can be varied for each ion listed above. Our previous studies of Ca II in the Galactic halo (Richter et al. 2005, 2009; Ben Bekhti et al. 2008, 2011; Wakker et al. 2007, 2008) and in intervening absorbers at low redshift (Richter et al. 2011) have demonstrated that Ca II is an excellent tracer for the distribution of neutral and partly ionized gas and its velocity-component structure, even in regions where Ca II is not the dominant ionization state.

Based on the Ca II model, and using a modified version of *FITLYMAN*, we have calculated for each individual UV line a synthetic absorption profile, for which we have




FIG. 4.— COS velocity profiles of the intermediate and high ions Si III, C IV and Si IV. The velocity structure in these ions is very different from that of the low ions, suggesting that they trace spatially distinct regions within the MS. High-velocity Si IV absorption between +150 and +200 km s^{-1} in the Si IV $\lambda 1402.8$ line is possibly blended with intergalactic absorption.

convolved an initial Voigt profile with input parameters (v_i, b_i, N_i) for $i = 1 \dots 7$ with the COS line-spread function (LSF) that is appropriate for the wavelength of the line. For the COS LSF we have used the improved LSF model described by Kriss (2011). Then, the columns N_i were varied to minimize the differences between the synthetic absorption profile and the observed COS data. This method delivers reliable (total) column densities for those ions that have multiple transitions with substantially different oscillator strengths in the COS wavelength range and for individual lines that are not fully saturated. More details about the accuracy of this method are presented in Sect. 3.1 and in the Appendix.

For the intermediate and high ions (Si III, Si IV, C IV) the component structure (and b values) are expected to be different from that for the low ions (as the gas phase traced by these ions often is spatially distinct from the gas phase traced by the low ions), but no information is available on the true component structure of these ions from the optical data. Therefore, we did not try to model the absorption profiles of the high-ion lines, but estimated total column densities (and limits) for these ions solely from the AOD method. Similarly, we used solely the AOD method to determine the column density of C II*.

For the analysis of H₂ in the Magellanic Stream detected in the Fairall 9 *FUSE* data (Richter et al. 2001a), we have modeled the H₂ absorption using synthetic spectra generated with FITLYMAN. As model input we take

into account the component structure and line widths seen in the optical Ca II/Na I absorption, together with a Gaussian LSF according to the *FUSE* spectral resolution. H₂ wavelengths and oscillator strengths have been adopted from Abgrall & Roueff (1989). Details on the H₂ modeling are presented in Sect. 3.3.

3. COLUMN-DENSITY MEASUREMENTS

3.1. Metal absorption in the optical

Optical absorption related to gas in the Magellanic Stream in the velocity range 120–230 km s^{-1} (this velocity range is defined by the H I 21 cm data of the Stream) is detected in the UVES spectrum of Fairall 9 in the lines of Ca II ($\lambda\lambda 3934.8, 3969.6$), Na I ($\lambda\lambda 5891.6, 5897.6$), and Ti II ($\lambda 3384.7$; 3.4σ detection). Fig. 2 shows the normalized absorption profiles of these ions plotted on the LSR velocity scale together with the H I 21cm emission profile from GASS. Ca II absorption is detected in seven individual absorption components centered at $v_{\text{LSR}} = +143, +163, +172, +188, +194, +204$ and $+218 \text{ km s}^{-1}$ with column densities in the range $\log N(\text{Ca II}) = 11.29 - 11.68$. The red solid line in Fig. 2 indicates the best Voigt-profile fit to the data. All Ca II, Na I and Ti II column-density measurements are summarized in Table 1.

The strongest Ca II absorption component in the MS (in terms of the absorption depth) is component 4 at $+188 \text{ km s}^{-1}$; this very narrow component is also detected in both Na I lines (Fig. 2). From the simultaneous fit of the Ca II and Na I doublets we obtain a very small Doppler parameter for component 4 of $b = 1.8 \text{ km s}^{-1}$. The small b value for this component is confirmed by fits to the individual lines of Ca II and Na I, which all imply $b < 2 \text{ km s}^{-1}$. The detection of Na I together with the small b value indicates that the gas in component 4 is relatively cold and dense and possibly is confined in a dense core with little turbulence. Components 3, 4, and 5 are also detected in Ti II (at 3.4σ significance; Fig. 2). Since Ti II and H I have almost identical ionization potentials (Table 2), the detection of Ti II suggests that most of the neutral gas column density is contained in these three components. This conclusion is supported by the H I 21cm emission profile from GASS (Fig. 2; upper panel), which also has its maximum in the velocity range between +170 and +200 km s^{-1} . Weak and relatively broad ($b = 7.4 \text{ km s}^{-1}$) Na I absorption is also detected in component 1 at $+143 \text{ km s}^{-1}$.

As mentioned in Sect. 2.5, we also have used the AOD method to determine the total column densities for Ca II, Na I and Ti II by integrating over the velocity range relevant for MS absorption (i.e., $+130 \text{ km s}^{-1} \leq v_{\text{LSR}} \leq +230 \text{ km s}^{-1}$.) The column densities derived by these two different methods agree very well within their 1σ error ranges (see Table 1, rows 10 and 11).

3.2. Metal absorption in the UV

In Fig. 3 we show normalized UV absorption profiles of the low ions C II, N I, O I, Al II, Si II, S II, and Fe II. The absorption profiles of the intermediate and high ions Si III, C IV and Si IV are shown in Fig. 4. Following the procedure described in Sect. 2.5, we have reconstructed the absorption pattern of UV metal lines of the low ions, using the component model defined by the optical lines of

TABLE 1
SUMMARY OF ION COLUMN-DENSITY MEASUREMENTS

Component		1	2	3	4	5	6	7		
v [km s ⁻¹]		+143	+163	+172	+188	+194	+204	+218		
b [km s ⁻¹]		7.4	7.9	6.4	1.8	3.4	4.9	5.1		
σ_b [km s ⁻¹]		2.4	4.1	2.2	0.3	1.4	1.7	2.0		
Ion	Instrument	log N_1	log N_2	log N_3	log N_4	log N_5	log N_6	log N_7	log $N_{\text{tot,fit}}$	log $N_{\text{tot,AOD}}$
H I	GASS	...	...	...	...	...	...	...	...	19.95
Na I	VLT/UVES	10.92	...	...	11.05	...	...	...	11.29 ± 0.04	11.33 ± 0.05
Ca II	VLT/UVES	11.58	11.29	11.63	11.68	11.58	11.43	11.35	12.37 ± 0.02	12.38 ± 0.03
Ti II	VLT/UVES	...	...	11.11	11.15	11.24	...	...	11.27 ± 0.06	≤ 12.00
C I	HST/COS	...	...	...	...	...	...	...	...	≤ 13.02
C II	HST/COS	...	...	...	...	...	...	...	...	≥ 14.93
C II*	HST/COS	...	...	...	...	...	...	...	...	13.35 ± 0.07
C IV	HST/COS	...	...	...	...	...	...	...	...	13.73 ± 0.04
N I	HST/COS	13.65	13.42	13.84	14.10	13.98	13.54	13.46	14.63 ± 0.05	≥ 14.51
O I	HST/COS	...	...	...	...	...	...	...	...	≥ 14.77
Al II	HST/COS	12.55	12.53	12.92	12.58	12.69	12.37	12.24	13.45 ± 0.06	≥ 13.18
Si II	HST/COS	14.10	13.75	14.17	14.13	14.24	13.92	13.89	14.90 ± 0.04	15.01 ± 0.17
Si III	HST/COS	...	...	...	...	...	...	...	...	≥ 13.57
Si IV	HST/COS	...	...	...	...	...	...	...	...	13.03 ± 0.06
P II	HST/COS	...	...	...	...	...	...	...	...	≤ 12.87
S II	HST/COS	13.95	13.70	14.03	14.00	14.09	13.77	13.74	14.77 ± 0.02	14.77 ± 0.04
Fe II	HST/COS	13.47	13.61	13.94	13.85	13.94	13.67	13.65	14.61 ± 0.04	14.59 ± 0.04
Ni II	HST/COS	...	...	...	...	...	...	...	...	≤ 13.37

Ca II. As already mentioned, the modeling method provides relevant results only for those ions that have multiple transitions in the available COS wavelength band and for single lines that are not (or at most mildly) saturated. In our case, the modeling method could be used to determine column densities for N I, Al II, Si II, S II, and Fe II, while for the fully saturated (single) lines of C II and O I the modeling does not yield relevant column density limits. In Fig. 3, the best-fitting column-density model for each ion is overlaid with the red solid line. The best-fitting model assumes the same b values for the individual subcomponents as derived for Ca II from the Voigt fit of the UVES data. The ion column densities for each velocity component are listed in Table 1 in rows 2 – 7; the total column density for each ion is given in row 10. As can be seen in Fig. 3, the absorber model successfully reproduces the shape of the absorption lines in the UV.

From Table 1 follows that the relative column-densities in the seven velocity components differ slightly from ion to ion. These differences are expected, because most ions have different ionization potentials and therefore trace slightly different gas phases within the absorber. In addition, differential dust depletion for some elements may also be relevant in this context. Because of the relatively limited spectral resolution of the COS instrument and the resulting overlap between neighbouring velocity components the accuracy for the determination of the column densities for individual subcomponents is relatively low (~ 0.30 dex, typically). Much better constrained are the total column densities (Table 1, row 10) for the ion multiplets N I, S II, Si II and Fe II, which have 1σ uncertainties < 0.05 dex. This is because the integrated (total) column density for each of these ions is determined predominantly by the integrated equivalent widths of the weaker lines and the equivalent width *ratios* of the weaker and stronger transitions. In the Appendix we present additional model plots to provide a more detailed insight into

the allowed parameter range (b_i, N_i) for the individual subcomponents and their impact on the shape of the absorption profiles and total column-density estimates.

Next to the absorption modeling we have used the AOD method to determine total column densities (or column-density limits) in the COS data for each of the ions listed above (Table 1, row 11). For the unsaturated lines of S II, Si II, and Fe II the total column densities derived from the AOD method are in excellent agreement ($< 1\sigma$) with the total column densities determined from the profile modeling. Note that the absorption of C II* in the MS will be discussed separately in Sect. 4.4.

3.3. H I 21cm emission

The 21cm velocity profile shown in the upper panel of Fig. 2 indicates H I emission from gas in the Magellanic Stream in the LSR velocity range between +100 and +250 km s⁻¹. The MS emission shows an asymmetric pattern with a peak near $v_{\text{LSR}} \approx +180$ km s⁻¹, surrounded by weaker satellite components at lower and higher radial velocities. The overall shape of the emission profile mimics that of the inversed absorption profiles of the low-ion lines of N I and S II indicating that both the pencil-beam absorption data and the GASS emission data, that have a beam-size of 15'6, sample the same physical structure. Integrating the GASS data in the above given velocity range for the single pixel that covers the position of Fairall 9 we obtain a total H I column density in the Stream of $\log N(\text{H I}) = 19.95$ ($N(\text{H I}) = (8.96 \pm 0.09) \times 10^{19}$ cm⁻²). The pixel value is somewhat smaller than the beam-averaged column density, which is $\log N(\text{H I}) = 19.98$. For comparison, Gibson et al. (2000) find $\log N(\text{H I}) = 19.97$ ($N(\text{H I}) = (9.35 \pm 0.47) \times 10^{19}$ cm⁻²) using Parkes 21cm observation of Fairall 9 with a beam size of 14'0. These values agree with each other within their 1σ error ranges. From the 21cm data of the *Leiden-Argentine-Bonn Survey* (LAB; Kalberla et

al. 2005) with ~ 0.5 deg resolution we obtain for the same velocity range an H I column density in the Stream of $\log N(\text{H I}) = 19.97$, thus very similar.

For the determination of metal abundances (using the GASS data in combination with the pencil-beam UVES and COS data) we adopt the value of $\log N(\text{H I}) = 19.95$ for the total H I column density in the Stream toward Fairall 9, together with an appropriate estimate of the systematic error of 0.03 dex that accounts for the limited spatial resolution of the 21cm data. As it is shown in the Appendix, this H I column density in the MS is in agreement with the observed shape of the Ly α absorption line in the COS data of Fairall 9, which provides an independent (albeit less stringent) measure for the neutral-gas column density in the Stream in this direction (see also Wakker, Lockman, & Brown 2011 for a detailed discussion on this topic).

3.4. Molecular hydrogen absorption

The detection of H_2 absorption in the Magellanic Stream in the *FUSE* data of Fairall 9 has been reported previously by Richter et al. (2001a) and Wakker (2006). H_2 absorption is seen near $v_{\text{LSR}} = +190 \text{ km s}^{-1}$ in the rotational levels $J = 0, 1$ and 2. Although the overall S/N in the *FUSE* data is low (≤ 15 per resolution element; see Sect. 2.5), the H_2 absorption is clearly visible in many lines due to the relatively high H_2 column density (Richter et al. 2001a).

We have reanalyzed the Fairall 9 *FUSE* data based on an improved data reduction pipeline and the component model discussed in the previous sections. While Richter et al. (2001) have used a curve-of-growth technique to derive the column densities $N(J)$ and b values of the H_2 rotational states $J = 0, 1, 2$, assuming as single MS absorption component, we here model the absorption spectrum of the MS H_2 absorption in the *FUSE* Fairall 9 data using synthetic spectra. For our model we assume that the H_2 absorption arises in the region with the highest gas density in the Stream, which is component 4 at $v_{\text{LSR}} = +188 \text{ km s}^{-1}$ detected in the UVES data in Ca II, Na I, and Ti II (Table 2). In fact, from absorption studies of Na I and H_2 in interstellar gas it is well known that these two species trace the same gas phase (i.e., cold neutral and cold molecular gas; e.g., Welty et al. 2006).

We adopt $b_{\text{turb}} = 1.8 \text{ km s}^{-1}$ for the H_2 absorption in component 4, as suggested by the Na I and Ca II absorption in the same component (Table 2). We further assume (and prove later) that the kinetic temperature of the H_2 absorbing gas is $T_{\text{kin}} \approx 100 \text{ K}$ in component 4. It is usually assumed that the Doppler parameter is composed of a turbulent and a thermal component, so that $b^2 = b_{\text{turb}}^2 + b_{\text{th}}^2$. For light elements and molecules (such as H_2) b_{th} must not be neglected. We therefore adopt $b(\text{H}_2) = 2.0 \text{ km s}^{-1}$ for our H_2 model spectrum.

Fig. 4 shows our best fitting H_2 model (red solid line) together with the *FUSE* data for the wavelength range between 1070 and 1080 Å. Since the b value is fixed to $b(\text{H}_2) = 2.0 \text{ km s}^{-1}$ and the velocity to $v_{\text{LSR}} = +188 \text{ km s}^{-1}$, the only free parameters in our model are the H_2 column densities $N(J)$. Table 2 summarizes the best fitting H_2 column densities using this method. The given errors represent the uncertainties for $\log N(J)$ under the assumption that $b = 2.0 \text{ km s}^{-1}$. The result-

TABLE 2
SUMMARY OF H_2 COLUMN-DENSITY MODELING

J	b^a [km s^{-1}]	$\log N(J)$	T_J [K] b^b
0	2.0	17.53 ± 0.24	93^{+149}_{-39}
1	2.0	17.68 ± 0.26	93^{+149}_{-39}
2	2.0	16.46 ± 0.21	< 172
3	2.0	≤ 15.60	< 172
Sum		$17.93^{+0.19}_{-0.16}$	

^a b -value fixed to 2.0 km s^{-1} ; see Sect. 3.4.

^b Equivalent Boltzmann temperature implied by $N(J)$; see Sect. 4.4.2.

ing total H_2 column density is $\log N(\text{H}_2) = 17.93^{+0.19}_{-0.16}$ and the fraction of hydrogen in molecular form is $\log f_{\text{H}_2} = \log [2N(\text{H}_2)/(N(\text{H I}) + 2N(\text{H}_2))] = -2.03$. Note that the value for $\log f_{\text{H}_2}$ derived in this way represents a sightline-average; the local molecular fraction in the H_2 absorbing region must be higher (because $N(\text{H I})$ in this region is smaller).

The H_2 column densities we derive from our spectral modeling are substantially (~ 1 dex) higher than the values presented in the earlier studies of the Fairall 9 *FUSE* data (Richter et al. 2001a; Wakker 2006). The reason for this discrepancy is the substantially lower b value that we adopt here. This more realistic b value for the H_2 absorbing gas is a direct consequence of the previously unknown complex absorption pattern in the MS toward Fairall 9, that is visible only in the high-resolution optical data of Ca II and Na I (see Sect. 3.1). Our study suggests, that, without the support of high-resolution and high S/N optical or UV data, that can be used to resolve the true sub-component structure of the absorbers and to deliver reliable b values for the subcomponents, the interpretation of interstellar H_2 absorption lines in low-resolution and low-S/N data from *FUSE* can be afflicted with large systematic uncertainties.

4. PHYSICAL AND CHEMICAL CONDITIONS IN THE GAS

4.1. Ionization conditions and overall metallicity

The presence of high, intermediate and low ions, as well as molecular hydrogen, in the Magellanic Stream toward Fairall 9 in seven individual velocity components implies a complex multi-phase nature of the gas. To obtain meaningful results for the chemical composition and physical conditions from the measured ion column densities, it is necessary to consider in detail the ionization conditions in the absorbing gas structure. For an element M we define its relative gas-phase abundance compared to the solar abundance in the standard way $[\text{M}/\text{H}] = \log (\text{M}/\text{H}) - \log (\text{M}/\text{H})_{\odot}$. Solar reference abundances have been adopted from Asplund et al. (2009).

The best ion to study the overall metallicity of neutral and weakly ionized interstellar gas is O I. Neutral oxygen and neutral hydrogen have similar ionization potentials and there is a strong charge-exchange reaction that couples both ions in gas with sufficiently high density. For our sightline, however, the O I column density is not well constrained because of the saturation of the O I $\lambda 1302.2$ line and the lack of other, weaker O I transitions in the COS wavelength range. Because it is impossible to measure the O I column density in the MS using our




FIG. 5.— Continuum-normalized *FUSE* spectrum of Fairall 9 in the wavelength range between 1075 and 1081 Å. Absorption by H_2 from the rotational levels $J = 0, 1$ and 2 related to gas in the Magellanic Stream are indicated above the spectrum using the common notation scheme. The best-fitting model for the H_2 absorption is overplotted with the red solid line.

component models and the AOD method to a satisfactory accuracy, we do not further consider the $\text{O I } \lambda 1302.2$ line for our analysis.

For the determination of the overall metallicity of the gas in the Magellanic Stream toward Fairall 9 we choose instead S II , which is another useful ion for measuring interstellar abundances, as singly-ionized sulfur is an excellent tracer for neutral hydrogen without being depleted into dust grains (e.g., Savage & Sembach 1996). We also have a very accurate column-density determination of $\log N(\text{S II}) = 14.77 \pm 0.02$ in the Magellanic Stream, based on the absorption modeling of the two S II lines at 1250.6 and 1253.8 Å and (independently) from the AOD method (see Table 1). Our value is slightly higher than the S II column density of $\log N(\text{S II}) = 14.69 \pm 0.03$ derived by Gibson et al. (2000) from *HST*/GHRS data.

From the measured ($\text{S II}/\text{H I}$) ratio we obtain an initial estimate of the sulfur abundance in the Stream toward Fairall 9 of $[\text{S}/\text{H}] \approx -0.30$. Since the ionization potential of S II (23.2 eV; see Table 3) is higher than that of H I (13.6 eV), we have to model the ionization conditions in the gas to obtain a more precise estimate for the sulfur abundance (and the abundances of the other detected metals) in the Stream toward Fairall 9. While previous studies of Galactic halo clouds (Wakker et al. 1999; Gibson et al. 2000) have demonstrated that the ($\text{S II}/\text{H I}$) ratio is a very robust measure for the sulfur-to-hydrogen ratio, hardly being affected by the local ionizing radiation field, the ionization modeling is important to obtain a reliable estimate of the systematic uncertainty that arises from the use of S II as reference indicator for the overall α abundance in the gas. For this task, we have used the photoionization code *Cloudy* (v10.00; Ferland et al. 1998), which calculates the expected column densities for different ions as a function of the ionization parameter $U = n_\gamma/n_{\text{H}}$ for a gas slab with a given neutral gas column density and metallicity, assuming that the slab is illuminated by an external radiation field. As radiation field we adopt a combined Milky Way plus extragalactic ionizing radiation field based on the models by Fox et al. (2005) and Bland-Hawthorn & Maloney (1999, 2002), appropriately adjusted to the position of the MS relative to the Milky Way disk.

While our absorption modeling provides rough estimates for the ion column densities in the seven sub-

components, we refrain from trying to model the ionization conditions in the individual components. This is because the unknown H I column densities in these components, together with the unknown geometry of the overall gas structure, would lead to large systematic uncertainties for such models. Instead, we have used *Cloudy* to obtain *integrated* elemental abundances in the gas of the Magellanic Stream toward Fairall 9 (representing the optical-depth weighted mean of the individual element abundances in all subcomponents). It is, however, important to emphasize that meaningful results for integrated elemental abundances from *Cloudy* can be obtained only for those ions for which the column densities in the ionization model do not strongly depend on the gas density (and ionization parameter), because the latter quantities are expected to vary *substantially* among the individual subcomponents.

For our *Cloudy* model we assume a neutral gas column density of $\log N(\text{H I}) = 19.95$, an overall metallicity of $[\text{Z}/\text{H}] = -0.3$, and solar *relative* abundances of the metals from Asplund et al. (2009), based on our results described above. In Fig. 5, upper panel, we display the expected logarithmic column densities for low and intermediate ions as a function of $\log U$ and $\log n_{\text{H}}$ for this model. The measured total column densities for these ions (Table 1, 10th column) are plotted on the right hand side of the panel. The total Ca II column density of $\log N(\text{Ca II}) = 12.37$ sets a limit for the (averaged) ionization parameter and density in the gas of $\log \langle U \rangle \leq -3.55$ and $\log \langle n_{\text{H}} \rangle \geq -1.95$. The corresponding density range that we consider as relevant for our *Cloudy* model ($2.5 \leq \log n_{\text{H}} \leq -1.95$) is indicated in Fig. 5 with the gray-shaded area. For this range, the expected column densities of N I , Si II , Fe II , S II , Al II , Ni II , and Ti II are nearly independent of $\log U$ and $\log n_{\text{H}}$. For S II , in particular, the column density varies within only 0.03 dex in the above given density range, supporting the previous conclusions from Wakker et al. (1999) and Gibson et al. (2000). In the lower panel of Fig. 5 we show the differences (ΔN) between the measured column densities and the mean column densities predicted by *Cloudy*. For Ni II and Ca II , the arrows indicate the upper limit and the allowed range for ΔN , respectively.

From the comparison between the predicted and measured ion column densities, together with the solar ref-




FIG. 6.— *Upper panel:* Cloudy photoionization model of low-ionization species in the MS toward Fairall 9. The colored lines indicate the predicted ion column densities as a function of the ionization parameter U and gas density n_H for a gas slab with a (total) neutral hydrogen column density of $\log N(\text{H I}) = 19.95$ and a metallicity of $[Z/H] = -0.30$. Observed ion column densities are indicated on the right-hand side with the filled colored circles. The gray-shaded area marks the range for U and n_H that is relevant for our study based on the observed Ca II column density (see Sect. 4.1). *Lower panel:* differences between the observed and the predicted ion column densities, $\Delta N = N_{\text{Cloudy}} - N_{\text{Obs.}}$; these differences are discussed in Sect. 4.1.

erence abundances from Asplund et al. (2009), we obtain the following gas-phase abundances: $[\text{N}/\text{H}] = -1.15 \pm 0.06$, $[\text{Si}/\text{H}] = -0.57 \pm 0.06$, $[\text{Fe}/\text{H}] = -0.86 \pm 0.06$, $[\text{S}/\text{H}] = -0.30 \pm 0.04$, $[\text{Al}/\text{H}] = -0.92 \pm 0.08$, $[\text{Ti}/\text{H}] = -1.62 \pm 0.07$, $[\text{Ni}/\text{H}] \leq -0.81$, and $[\text{Ca}/\text{H}] \leq 0$. The listed errors include the uncertainties from the column density measurements for the metal ions and H I (Table 1), the systematic uncertainty due to the beam size of the 21cm measurements (Sect. 3.3), and the systematic uncertainty for the ionization correction from the Cloudy model (see above). We here do not consider the systematic errors that come with the solar reference abundances (Asplund et al. 2009; see Table 3, third row), because these errors are irrelevant for the comparison between our results and other abundance measurements, if the identical reference values are used. All gas-phase abundances derived with Cloudy are listed in Table 3, sixth row.

The measured S/H ratio of $[\text{S}/\text{H}] = -0.30 \pm 0.04$ corresponds to an overall sulfur abundance in the Magellanic Stream toward Fairall 9 of $0.50^{+0.05}_{-0.04}$ solar. The value for $[\text{S}/\text{H}]$ is 0.25 dex higher than the value derived for the MS toward Fairall 9 by Gibson et al. (2000) based on GHRS data. Several factors contribute to this discrepancy: (a)

the somewhat higher (+0.08 dex) S II column density that we derive from the COS data, (b) the slightly lower H I column density (−0.02 dex) that we obtain from the GASS data, and (c) the substantially higher (0.15 dex) solar reference abundance of sulfur (Asplund et al. 2009) that we use to derive $[\text{S}/\text{H}]$ compared to the value from Anders & Grevesse (1989) used by Gibson et al. (2000).

Remarkably, the sulfur abundance we derive here is *higher* than the present-day stellar sulfur abundances in the SMC ($[\text{S}/\text{H}] = -0.53 \pm 0.15$) and in the LMC ($[\text{S}/\text{H}] = -0.42 \pm 0.09$; Russell & Dopita 1992), but matches the present-day interstellar S abundances found in H II regions in the Magellanic Clouds (Russell & Dopita 1990). This interesting result will be further discussed in Sect. 6. Because of the large abundance scatter of sulfur in the Magellanic Clouds and because S appears to be underabundant compared to oxygen in the LMC and overabundant compared to oxygen in the SMC (Russell & Dopita 1992), the measured (S/H) ratio does not provide a constraining parameter to pinpoint the origin of the gaseous material in the Stream toward Fairall 9 in either SMC or LMC.

From our Cloudy model it follows that all other elements listed above have gas-phase abundances (or abundance limits) that are lower than that of S. Next to the intrinsic nucleosynthetic MS abundance pattern, dust depletion is another important effect that contributes to deficiency of certain elements (e.g., Ti, Al, Fe, Si) in the gas phase. This aspect will be discussed in Sect. 4.3. Note, again, that we do not attempt to constrain a single value for $\log U$ or $\log n_H$ with our Cloudy model, as our sightline passes a complex multiphase structure that is expected to span a substantial *range* in gas densities and ionization parameters.

In this paper, we do not further analyze the gas phase that is traced by the intermediate and high ions Si III , C IV and Si IV , which probably is tracing the ionized boundary layer between the neutral gas body of the Stream and the ambient hot coronal halo gas of the Milky Way (see Fox et al. 2010). For completeness, we show in Fig. 4 the velocity profiles of the available transitions of Si III , C IV and Si IV in the COS spectrum of Fairall 9. The velocity structure in these ions clearly is different from that of the low ions, suggesting that they trace spatially distinct regions in the MS. The high-ion absorption toward Fairall 9 will be included in a forthcoming paper that will discuss those ions in several lines of sight intersecting the Magellanic Stream (A.J. Fox et al. 2014, in preparation).

4.2. Nitrogen abundance

The measured gas phase abundance of nitrogen in the Magellanic Stream toward Fairall 9 is $[\text{N}/\text{H}] = -1.15 \pm 0.06$, which is 0.85 dex lower than the abundance of sulfur. In diffuse neutral gas the depletion of nitrogen into dust grains is expected to be very small (Savage & Sembach 1996). Consequently, the low nitrogen abundance in the Stream reflects the nucleosynthetic enrichment history of the gas and thus provides important information about the origin of the gas.

For the SMC, Russell & Dopita (1992) derive a mean present-day stellar nitrogen abundance of $[\text{N}/\text{H}] = -1.20$. This value is close to the value derived for the MS toward Fairall 9, although the gas in Stream has not

TABLE 3
SUMMARY OF INTEGRATED GAS-PHASE ABUNDANCES AND DUST DEPLETION VALUES

Ion (X)	Z	I.P. ^a [eV]	$\log (M/H)_{\odot}^{b}$ +12	$\log (X/H\text{I})^c$	$[M/H]^d$	$\log \delta(M)$
C I	6	11.26	8.43 ± 0.05	≤ -6.93	...	...
C II	6	24.38	8.43 ± 0.05	≥ -5.02	...	...
N I	7	14.53	7.83 ± 0.05	-5.32 ± 0.05	-1.15 ± 0.06	...
O I	8	13.62	8.69 ± 0.05	≥ -1.87	...	...
Na I	11	5.14	6.24 ± 0.04	$-8.66 \pm 0.0x$	...	...
Al II	13	18.83	6.45 ± 0.03	-6.50 ± 0.06	-0.92 ± 0.08	-0.62
Si II	14	16.35	7.51 ± 0.03	-5.05 ± 0.04	-0.57 ± 0.06	-0.27
P II	15	19.77	5.41 ± 0.04	≤ -7.08	...	...
S II	16	23.34	7.12 ± 0.03	-5.18 ± 0.02	-0.30 ± 0.04	(0)
Ca II	20	11.87	6.34 ± 0.04	-7.58 ± 0.02	≤ -0.30	0 to -1.58
Ti II	22	13.58	4.95 ± 0.05	-8.68 ± 0.06	-1.63 ± 0.07	-1.33
Fe II	26	16.12	7.50 ± 0.04	-5.34 ± 0.04	-0.86 ± 0.06	-0.56
Ni II	28	18.17	6.22 ± 0.04	≤ -6.58	≤ -0.81	≤ -0.51

^a I.P.= ionization potential.

^b Solar reference abundance for element M from Asplund et al. (2009).

^c Listed errors include uncertainties from the column-density measurements of X and H I.

^d Gas phase abundance for element M, defined as $[M/H] = \log (M/H) - \log (M/H)_{\odot}$, derived from the CLOUDY model. The listed errors include uncertainties from the column-density measurements, the beam-smearing of the 21cm data, and the ionization correction from the Cloudy modeling.

^e Depletion value, defined as $\log \delta(M) = [M/H]_{\text{MS}} - [S/H]_{\text{MS}}$.

been enriched with nitrogen since it was stripped off its parent galaxy. The nitrogen abundances found in SMC H II regions and supernova remnants span a rather wide range of $-1.37 \leq [N/H] \leq -0.95$ (Russell & Dopita 1990). Thus, the observed nitrogen abundance in the MS toward Fairall 9 would be in accordance with an SMC origin of the gas only, if the (mean) nitrogen abundance in the SMC has not increased substantially after the Stream was separated $\sim 1 - 2$ Gyr ago.

In the LMC, in contrast, the mean present-day stellar nitrogen abundance is found to be much higher ($[N/H] = -0.69$). The N abundance range in LMC H II regions and supernova remnants is $[N/H] = -0.86$ to -0.38 , the minimum $[N/H]$ still being substantially higher than the nitrogen abundance in the Stream toward Fairall 9. Therefore, an LMC origin of the MS is plausible only if the LMC has substantially increased its nitrogen abundance during the last ~ 2 Gyr.

In Sect. 5.1 we will further consider these interesting aspects and discuss the observed N abundance in the MS in the context of the enrichment history of the Magellanic Clouds.

4.3. Dust depletion pattern

Heavy elements such as Al, Si, Fe, Ca, Ti, Fe and Ni are known to be strongly depleted into dust grains in interstellar gas in the Milky Way and other galaxies, while other elements such as S, O, N are only very mildly or not at all affected by dust depletion (e.g., Savage & Sembach 1996; Welty et al. 1997).

To study the depletion pattern of these elements in the Magellanic Stream we define the depletion values from the relative abundance of each depleted element, M, in relation to the abundance of sulfur. We define accordingly $\log \delta(M) = [M/H]_{\text{MS}} - [S/H]_{\text{MS}}$. Using this definition, the depletions values are $\log \delta(\text{Si}) = -0.27$, $\log \delta(\text{Fe}) = -0.56$, $\log \delta(\text{Ti}) = -1.32$, $\log \delta(\text{Al}) = -0.62$, and $\log \delta(\text{Ni}) \leq -0.51$ (see also Table 3, seventh row). Such depletion values are typical for warm Galactic halo

clouds, as derived from UV absorption-line measurements (see Savage & Sembach 1996; their Fig. 6).

Note that the depletion values (or limits) would be different, if the *relative* chemical abundances of the depleted elements would differ from the relative chemical composition of the sun (Asplund et al. 2009). In view of the abundance patterns found in the LMC and SMC (Russell & Dopita 1990, 1992), this is actually a likely scenario, but no further conclusions can be drawn at this point without knowing the exact intrinsic chemical composition of the MS gas toward Fairall 9.

Since the expected Ca II column density in MS gas is expected to depend strongly on the local gas density and ionization parameter (Fig. 6), the depletion value of Ca cannot be tightly constrained, but may lie anywhere in the range $\log \delta(\text{Ca}) \geq -1.58$. Under typical interstellar conditions, Ca is strongly depleted into dust grains with large depletion values that are similar to those of other elements that have condensation temperatures above $T = 1500$ K (e.g., Ti; Savage & Sembach 1996). If one *assumes* that $\log \delta(\text{Ca}) \approx \log \delta(\text{Ti})$ for the MS toward Fairall 9, then Fig. 6 would imply that the average density in the neutral gas in the Stream in this direction is $\log n_{\text{H}} \approx 0$ or $n_{\text{H}} \approx 1 \text{ cm}^{-3}$. This density, together with the neutral gas column density, would imply a characteristic thickness of the absorbing neutral gas layer of $d = N(\text{H I})/n_{\text{H}} \approx 30 \text{ pc}$.

At first glance, one may regard this as a relatively small value. The complex velocity-component structure of the absorber indicates, however, that the absorber is composed of several small cloudlets along the line of sight that (together) are extending over a spatial range that is larger than the value indicated by the *mean* thickness. In view of the relatively high metal abundance derived for the kinematically complex absorber toward Fairall 9, the presence of a population of metal-enriched gas clumps at small scales in the MS is not an unlikely scenario.




FIG. 7.— COS velocity profiles of C II* $\lambda 1335.7$ (upper panel). C II* absorption from the Magellanic Stream is visible between +120 and +200 km s^{-1} . For comparison, the velocity profile of S II $\lambda 1250.6$ is also shown (lower panel). The component structure of the low ions is indicated with the tic marks.

4.4. Physical conditions in the H_2 absorbing region

As discussed in Sect. 3.1, the UVES spectrum of Fairall 9 shows the presence of Na I, Ca II, and possibly Ti II in component 4. In the same component, the detected H_2 absorption is expected to arise (Sect. 3.3), indicating the presence of a cold, dense (and predominantly neutral) gas phase in the MS in this direction. In the following, we want to combine the information from the different instruments to explore the physical and chemical conditions in the H_2 absorbing gas phase in the MS toward Fairall 9.

4.4.1. Gas density

While our Cloudy model does not provide any relevant information on the *local* gas densities in the individual subcomponents, we can use the observed molecular hydrogen fraction in the gas to constrain n_{H} in the H_2 absorbing component 4, assuming that the abundance of H_2 relative to H I is governed by a formation-dissociation equilibrium. In a formation-dissociation equilibrium the neutral to molecular hydrogen column density ratio in an interstellar gas cloud is given by

$$\frac{N(\text{H I})}{N(\text{H}_2)} = \frac{\langle k \rangle \beta}{R n(\text{H I}) \phi}, \quad (1)$$

where $\langle k \rangle \approx 0.11$ is the probability that the molecule is dissociated after photo absorption, β is the photo-absorption rate per second within the cloud, and R is the H_2 formation rate on dust grains in units $\text{cm}^3 \text{s}^{-1}$. For low molecular hydrogen fractions we can write $n_{\text{H}} = n(\text{H I}) + 2n(\text{H}_2) \approx n(\text{H I})$. The parameter $\phi \leq 1$ in equation (1) describes the column-density fraction of the H I that is physically related to the H_2 absorbing gas, i.e., the fraction of the neutral gas atoms that can be transformed into H_2 molecules (see Richter et al. 2003 for details). Our absorption modeling of the undepleted low ions S II and N I indicates that component 4 contains $\sim 15 - 30$ percent of neutral gas column, so that we assume $\phi = 0.15 - 0.30$ in lack of any more precise information.

For interstellar clouds that are optically thick in H_2 (i.e., for $\log N(\text{H}_2) \gg 14$) H_2 line self-shielding needs to be considered in equation (1), because the self-shielding




FIG. 8.— Rotational excitation of H_2 in the Magellanic Stream. The fit for the excitation temperatures (assuming a Boltzmann distribution) for $J = 0, 1$ and $J = 2, 3$ are indicated with the solid and dashed line, respectively.

reduces the photo-absorption rate (β) in the cloud interior. Draine & Bertoldi (1996) find that the H_2 self-shielding can be expressed by the relation $\beta = S \beta_0$, where $S = (N_{\text{H}_2}/10^{14} \text{cm}^{-2})^{-0.75} < 1$ is the self-shielding factor and β_0 is the photo absorption rate at the edge of the cloud. The parameter β_0 is directly related to the intensity of the ambient UV radiation field at the edge of the H_2 -bearing clump. Compared to the Milky Way disk, where UV bright stars dominate the mean photo-absorption rate of $\beta_{0,\text{MW}} = 5.0 \times 10^{-10} \text{s}^{-1}$ (e.g., Spitzer 1978), the value for β_0 in the Magellanic Stream is expected to be substantially smaller. This is because the solid angle of the Milky Way disk is relatively small at 50 kpc distance and the contribution of the extragalactic UV background to $\beta_{0,\text{MS}}$ is expected to be (relatively) small, too. From the model by Fox et al. (2005) follows that the UV flux between 900 and 1100 $\text{\AA}$ is reduced by a factor of 160 compared to the mean flux within the Milky Way disk, so that we assume $\beta_{0,\text{MS}} = 3.1 \times 10^{-12} \text{s}^{-1}$. With a total H_2 column density of $8.5 \times 10^{17} \text{cm}^{-2}$ in the Stream the self-shielding factor S becomes 1.13×10^{-3} , so that the photo-absorption rate in the cloud core is estimated as $\beta = 3.5 \times 10^{-15} \text{s}^{-1}$. For the H_2 grain formation rate in the Magellanic Stream we adopt the value derived for the SMC based on *FUSE* H_2 absorption-line data (Tumlinson et al. 2002), i.e., $R_{\text{MS}} = 3 \times 10^{-18} \text{cm}^3 \text{s}^{-1}$. This value is 10 times smaller than R in the disk of the Milky Way (Spitzer 1978).

If we solve equation 3 for $\phi n_{\text{H}} = \phi n(\text{H I})$ and include the values given above, we obtain a density of $\phi n_{\text{H}} \approx 1 \text{cm}^{-3}$ or $n_{\text{H}} \approx 4 - 8 \text{cm}^{-3}$ for $\phi = 0.15 - 0.30$. These densities are very close to the density derived for the H_2 absorbing gas in the Leading Arm of the MS toward NGC 3783 ($n_{\text{H}} \approx 10 \text{cm}^{-3}$; Sembach et al. 2001). Note that H_2 absorption has also been detected in the Magellanic Bridge (Lehner 2002).

4.4.2. Gas temperature

In Fig. 8 we plot the measured logarithmic H_2 column density for each rotational state, $\log N(J)$, divided by its statistical weight, g_J , against the rotational excitation energy, E_J . Rotational excitation energies and statistical weights have been adopted from the compilation of Morton & Dinerstein (1976). The data points in Fig. 8 can be fitted by a Boltzmann distribution (i.e., a straight line in this plot), where the slope characterizes the (equivalent) excitation temperature for the rotational levels considered for the fit. For the two rotational ground states $J = 0, 1$ the Boltzmann distribution thus is given by the equation $N(1)/N(0) = g_1/g_0 \exp(-E_{01}/kT_{01})$. Using this equation, together with our measured column densities for $J = 0, 1$, we find an excitation temperature of $T_{01} = 93^{+149}_{-39}$ K (Fig. 8 solid line). Because the two rotational ground states most likely are collisionally excited, T_{01} represents a robust measure for the kinetic temperature in the H_2 absorbing gas (e.g., Spitzer 1978); however, because of the saturation of the H_2 lines and the resulting relatively large errors for $N(0, 1)$, the uncertainty for T_{01} is substantial. In a similar manner, we derive for the rotational states $J = 2, 3$ an upper limit for the equivalent Boltzmann temperature of $T_{23} \leq 172$ K (Fig. 8 dashed line). This higher excitation temperature for $J \geq 2$ reflects excitation processes other than collisions, such as UV pumping and H_2 formation pumping (Spitzer 1978). However, the relatively low temperature limit of $T_{23} \leq 172$ K reflects the low intensity of the ambient dissociating UV radiation field at the position of the MS.

The above given excitation temperatures deviate from our earlier estimates (Richter et al. 2001a). This is not surprising, however, since the values for $N(J)$ have substantially changed, too (see Sect. 3.3). The derived value of $T_{01} = 93$ K lies in a temperature regime that is typical for diffuse H_2 gas in the disk of the Milky and in the Magellanic Clouds (e.g., Savage et al. 1977; Tumlinson et al. 2002; de Boer et al. 1998; Richter et al. 1998; Richter 2000).

From the density and temperature estimate in the H_2 absorbing gas we now are able to provide a direct estimate of the thermal pressure, $P/k = nT$, in the cold neutral gas in the MS. With $n_{\text{H}} = 8 \text{ cm}^{-3}$ and $T = 93$ K we obtain a pressure of $P/k = 744 \text{ cm}^{-3} \text{ K}$. This value is in good agreement with previous estimates for the thermal pressure in the cold neutral phase of the MS from H I 21cm measurements (e.g., Wakker et al. 2002).

The thickness of the H_2 absorbing structure in the MS is $d_{\text{H}_2} = \phi N(\text{H I})/n_{\text{H}} \approx 0.6 - 1.2 \text{ pc}$. This small dimension of the absorbing gas clump explains the very small velocity dispersion ($b = 1.8 \text{ km s}^{-1}$; see Table 1) that is measured for this component.

4.5. Electron density and C^+ cooling rate from C II^* absorption

In Fig. 7 (upper panel) we show the LSR velocity profile of the $\text{C II}^* \lambda 1335.7$ line, based on the COS data. Weak absorption is visible at MS velocities, but the component structure is different from that of the other low ions (e.g., S II ; Fig. 7, lower panel). Because we could not identify any other (e.g., intergalactic) origin for this absorption feature, we assume that it is caused by C II^* in the Magellanic Stream. The C II^* feature is characterized

by a stronger and broader absorption peak near $+150 \text{ km s}^{-1}$, falling together with components 1 and 2 defined in Table 1, and another weak and narrow component that is seen near $+190 \text{ km s}^{-1}$, most likely associated with component 4, in which H_2 , Na I , and Ca II is detected. Therefore, the relative strengths of the C II^* components appear to be inverted from those of the ground-state species. This indicates multi-phase gas, in which the ionization fractions and electron densities vary among the different absorption components.

Under typical interstellar conditions, the relative population of the fine-structure levels of ionized carbon (C^+) are governed by the balance between collisions with electrons and the radiative decay of the upper level into the ground state $2s^2 2p^2 P_{1/2}$. The $\text{C II}^* \lambda 1335.7$ transition arises from the $2s^2 2p^2 P_{3/2}$ state, which has an energy of $\sim 8 \times 10^8 \text{ eV}$ above the ground state. Measurements of the column-density ratios $N(\text{C II}^*)/N(\text{C II})$ thus can be used to estimate the electron densities n_e in different interstellar environments, including Galactic high-velocity clouds (HVCs) and circumgalactic gas structures (e.g., Zech et al. 2008; Jenkins et al. 2005).

For the C II^* absorption in the MS toward Fairall 9 we measure a total column density of $\log N(\text{C II}^*) = 13.35 \pm 0.07$ for the velocity range $v_{\text{LSR}} = 120 - 220 \text{ km s}^{-1}$ using the AOD method. For the weak absorption associated with component 4 we derive $\log N(\text{C II}^*) = 12.63 \pm 0.03$ (AOD). The $\text{C II} \lambda 1334.5$ absorption in the MS is fully saturated and thus does not provide a constraining limit on $N(\text{C II})$ (see Table 1). We therefore use S II as a proxy, because the $\text{S II}/\text{C II}$ ratio is expected to be constant over a large density range (Fig. 6) and both elements are not expected to be depleted into dust grains in the MS. Using a solar (S/C) ratio (Asplund et al. 2009), we estimate from the values listed in Table 1 that the total C II column density in the MS is $\log N_{\text{tot}}(\text{C II}) \approx 16.10$ and the column density in component 4 is $\log N_4(\text{C II}) \approx 15.30$.

Because the C^+ fine-structure population depends strongly on the gas temperature (see, e.g., Spitzer 1978), we do not attempt to estimate a mean value for $n(e)$ from $N_{\text{tot}}(\text{C II}^*)/N_{\text{tot}}(\text{C II})$, because we know that the temperature in this multi-phase absorber spans a large range between ~ 100 K and probably few 1000 K, so that the resulting estimate for $\langle n_e \rangle$ would be meaningless. Instead, we concentrate on component 4, for which we know the gas temperature and density from the analysis of the H_2 rotational excitation (Sect. 4.4). Keenan et al. (1986) have calculated electron excitation rates for the C^+ fine-structure transitions for a broad range of physical conditions in interstellar gas. Using our temperature and density estimates for the gas in component 4 ($T = 93$ K and $n_{\text{H}} = 4 - 8 \text{ cm}^{-3}$) together with their predicted population rate ratios ($2s^2 2p^2 P_{3/2}/2s^2 2p^2 P_{1/2}$) for $T = 100$ K and $n_{\text{H}} = 5 - 10 \text{ cm}^{-3}$ (their Fig. 2), the measured column-density ratio $\log[N(\text{C II}^*)/N(\text{C II})] = -2.67$ implies that the electron density in component 4 is small compared to n_{H} , namely $n_e \leq 0.05 \text{ cm}^{-3}$. This low value for n_e is in excellent agreement with expectations for a stable cold, neutral medium in gas with subsolar metallicities (Wolfire et al. 1995).

Having a robust measure for $N_{\text{tot}}(\text{C II}^*)$, it is also possible to estimate the C^+ cooling rate in the Magellanic Stream toward Fairall 9. Because ionized carbon is a ma-

major cooling agent of interstellar gas in a wide range of environments (e.g., Dalgarno & McCray 1972), the C^+ cooling rate is an important quantity that governs the thermal state of diffuse gas inside and outside of galaxies. For the hydrogen and electron densities in the Magellanic Stream (see above) the C^+ cooling rate is governed predominantly by spontaneous de-excitations, while collisional de-excitations can be neglected (e.g., Spitzer 1978). Following Lehner, Wakker & Savage (2004), the C^+ cooling rate per neutral hydrogen atom can be estimated as $l_C = 2.89 \times 10^{-20} N(C^+)/N(H\text{ I}) \text{ erg s}^{-1}$. If we adopt $\log N(H\text{ I}) = 19.95$ and $\log N(C\text{ II}^*) = 13.35$ we derive a mean (sightline average) C^+ cooling rate per neutral hydrogen atom of $\log l_C = -26.14$ for the MS toward Fairall 9. This value is almost one order of magnitude lower than the cooling rate derived for the one-tenth-solar metallicity HVC Complex C ($\log l_C \approx -27$), but is relatively close to the values derived for solar-metallicity clouds in the Milky Way disk and in the lower Galactic halo ($\log l_C \approx -26$; see Lehner, Wakker & Savage 2004, their Table 4). This result underlines the importance of metal cooling for the thermal state of metal-rich circumgalactic gas absorbers in the local Universe.

5. DISCUSSION

5.1. Enrichment history of the Magellanic Stream

5.1.1. Alpha and nitrogen abundances

One major result of our study is the surprisingly high metallicity of the MS toward Fairall 9. From the measured sulfur abundance of $[S/H] = -0.30 \pm 0.04$ follows that the α abundance in the Stream in this direction ($l = 295, b = -58$) is as high as 0.5 solar, which is ~ 5 times higher than the α abundance derived for other sightlines passing the MS toward NGC 7469 ($l = 347, b = -64$), RBS 144 ($l = 299, b = -66$), and NGC 7714 ($l = 88, b = -56$; Fox et al. 2010, 2013). Our detailed analysis of the ionization conditions in the gas, the comparison between H I 21cm measurements from different radio telescopes with different beam sizes, and the analysis of the H I Ly α absorption toward Fairall 9 (see Appendix) do not provide any evidence, that this high sulfur abundance could be a result from the various systematic errors that come along with our analysis (see also the discussion in Gibson et al. 2000 on this topic). We thus are forced to conclude that the measured high sulfur abundance in the gas reflects the true chemical composition of the Stream in this direction.

As a guide to the following discussion, we have plotted in Fig. 9 the derived MS abundances (black filled circles) together with the SMC and LMC present-day stellar and nebular abundances (Russell & Dopita 1992; Hughes et al. 1998; red (SMC) and green (LMC) filled circles). With a sulfur abundance of 0.5 solar, the MS toward Fairall 9 exhibits an α abundance that is *higher* than the average present-day α abundance in both SMC and LMC (~ 0.3 and ~ 0.4 solar, respectively; Russell & Dopita 1992). This finding has profound implications for our understanding of the origin of the Magellanic Stream and its enrichment history.

Early tidal models suggest that the MS was stripped from the Magellanic Clouds ~ 2 Gyr ago (Gardiner & Noguchi 1996). More recent models, that take into account that the Magellanic Clouds possibly are on their

first pass through the Milky Way halo, have included other relevant mechanisms that would help to unbind the gas, such as blowouts and tidal resonances (Connors et al. 2006; Nidever et al. 2008; Besla et al. 2010). One would expect that the Stream’s chemical composition must reflect that of its parent galaxy at the time, when the gas was separated from the Magellanic Clouds, since the MS does not contain stars that would increase its metallicity since then. Following the well-defined age-metallicity relations of SMC and LMC (Pagel & Tautvaišienė 1998; Harris & Zaritsky 2004; 2009), one would expect that the Stream’s present-day α abundance is at most ~ 0.2 solar if originating in the SMC and ~ 0.25 solar if originating in the LMC. While the observed MS abundances toward RBS 144, NGC 7469, and NGC 7714 are in line with these abundance limits (and actually favour an SMC origin for this part of the Stream; see Paper I), the sulfur abundance in the MS toward Fairall 9 is at least twice as high as expected from a simple tidal model that assumes a homogeneous pre-enrichment of the SMC/LMC gas (e.g., Dufour 1975) before the Stream was stripped off.

A second important finding of our study, that adds to this puzzle, is the relatively low nitrogen abundance in the MS toward Fairall 9. The $[N/S]$ ratio ($= [N/\alpha]$ ratio) is -0.85 dex, which is very low for the relatively high α abundance of $[\alpha/H] = -0.3$ when compared to nearby extragalactic H II regions and high-redshift damped Lyman α (DLA) systems (see Pettini et al. 2008, Jenkins et al. 2005, and references therein). The α -process elements O, S, and Si are believed to be produced by Type II supernovae from massive progenitor stars, while the production of N as part of the CNO cycle in stars of different masses is less simple (and not yet fully understood). The so-called “primary” nitrogen production occurs, when the seed elements C and O are produced within the star during the helium burning phase, while the for “secondary” nitrogen production these seed elements were already present when the star condensed out of the ISM (Pettini et al. 2002; Henry & Prochaska 2007). Primary N is believed to be produced predominantly by stars of intermediate masses on the asymptotic giant branch (e.g., Henry et al. 2000). Because of the longer lifetime of intermediate-mass stars, primary N therefore is expected to be released into the surrounding ISM with a time delay of ~ 250 Myr compared to α elements. At low metallicities less than ~ 0.5 solar (i.e., at LMC/SMC metallicities $\sim 1 - 2$ Gyr ago, when the MS was separated from its parent galaxy) the nitrogen production is predominantly primary (see Pettini et al. 2008).

The low $[N/S]$ and high $[S/H]$ ratios observed in the MS toward Fairall 9 therefore indicate an abundance pattern that is dominated by the α enrichment from massive stars and Type II supernovae, while only very little (primary) nitrogen was deposited into the gas.

5.1.2. Origin of the gas

The most plausible scenario that would explain such an enrichment history, is, that the gas that later became the Stream was *locally* enriched in the Magellanic Clouds $\sim 1 - 2$ Gyr ago with α elements by several supernova explosions in a star cluster or OB association, and then separated/stripped from the stellar body of the parent galaxy *before* the primary nitrogen was dumped into the




FIG. 9.— Comparison of present-day metal abundances in the SMC (blue dots), LMC (green dots), and in the Magellanic Stream toward Fairall9 (black dots). For SMC and LMC the metallicities are derived from stellar and nebular abundances (from Russell & Dopita 1992; Hughes et al. 1998).

gas and the metals could mix into the ambient interstellar gas. Eventually, the supernovae explosions may have pushed away the enriched material from the stellar disk so that the gas was already less gravitationally bound at the time it was stripped and incorporated into the Magellanic Stream. In this scenario, it is the present-day N abundance in the MS that defines the metallicity floor of the parent galaxy and thus provides clues to the origin of the gas.

Chemical evolution models of the Magellanic Clouds suggest that the mean metallicities of SMC and LMC 1–2 Gyr ago were 0.2–0.3 dex lower than today (Pagel & Tautvaisiene 1998; Harris & Zaritsky 2004, 2009). Therefore, and because of the substantially higher mean N abundance in the LMC compared to the SMC (see Fig. 9), the low N abundance measured in the Stream toward Fairall9 favours a SMC origin of the gas. An LMC origin cannot be ruled out, however, because the scatter in the present-day N abundances in the LMC is large (see Sect. 4.2) and the age-metallicity relations from Pagel & Tautvaisiene (1998) and Harris & Zaritsky (2004, 2009) may not apply to nitrogen because of its special production mechanism. In fact, considering the position of the Fairall9 sightline (see Paper I, Fig. 1) and the radial velocity of the MS absorption in this direction, the bifurcation model of Nidever et al. (2008) *predicts* that the high-velocity gaseous material towards Fairall9 is part of the LMC filament of the Stream and thus should have a different chemical composition than the SMC filament traced by the other MS sightlines (Paper I). Another aspect that may be of relevance in this context is the fact that the Fairall9 sightline lies only 14.3° on the sky from the SMC. The relative high metal-abundance in the Stream toward Fairall9 thus may reflect the increasing importance of continuous ram-pressure stripping of metal-enriched SMC gas as the MCs get closer to the Milky Way.

While the above outlined enrichment scenario appears feasible to explain qualitatively the observed trend seen in the abundance pattern in the MS toward Fairall9, the question arises, how many massive stars and subse-

quent supernova explosions would then be required to lift the [S/N] and [S/H] ratios to such a high level. To answer this question, one would first need to have an idea about volume and mass of the gas that is enriched in this manner. The fact that Fox et al. (2010, 2013) have determined a much lower α abundance in the Stream of ~ 0.1 solar toward three other MS sightlines using similar data possibly suggests, that the high metallicity and low [N/S] ratio toward Fairall9 represents a relatively local phenomenon.

In Sect. 4.3, we have estimated a thickness of the absorbing neutral gas layer in the Stream of $d \approx 30$ pc from the mean gas density ($n_H = 1.0 \text{ cm}^{-3}$) and the total column density, assuming that the dust depletion values of Ti and Ca are similar in the gas. In the following, we regard this value as a realistic *lower* limit for the true physical size of the absorbing gas region. If we adopt the SN yields from Kobayashi et al. (2006) and assume a spherical symmetry for the absorbing gas region with a diameter and density as given above, we calculate that only a handful of massive stars would be required to boost the α abundance from initially 0.1 solar to 0.5 solar in such a small volume of gas. A much higher number of high-mass stars would be required, however, if the volume to be enriched would be substantially larger, as we outline in the following. Because the total mass of a spherical gas cloud can be written as the product of cloud volume and gas density ($M \propto V n_H \propto d^3 n_H$), and because we use the relation $d = N_H / n_H$ to estimate d , we can write $M \propto n_H^2$ for a fixed (measured) hydrogen column density in the gas. Therefore, if the mean gas density in the MS toward Fairall9 would be 0.1 cm^{-3} rather than the assumed 1.0 cm^{-3} , then the diameter of the cloud would be 300 pc instead of 30 pc, but it would require 100 times more massive stars to enrich this large volume to the level required. In view of the stellar content of the most massive star forming regions in the Magellanic Clouds (e.g., 30 Doradus; Melnick 1985) and the possible presence of a major star burst in the MCs at the time, when the Magellanic Stream was formed (Harris & Zaritsky 2009; Weisz et al. 2013), a size of a few dozen up to a few hundred pc together with the proposed enrichment scenario is fully consistent with our understanding of the star-formation history of the Magellanic Clouds.

In summary, the observed abundance pattern in the MS toward Fairall9 suggests the presence of a high-metallicity gas filament in the Stream in this direction, possibly originating in a region with enhanced star-formation activity in the SMC or LMC $\sim 1-2$ Gyr ago. Our study indicates that the chemical composition and thus the enrichment history of the MS seems to be more complex than previously thought. The Stream possibly is composed of different gas layers that have different chemical compositions and that originate from different regions with local star-formation histories within the Magellanic Clouds.

Interestingly, the MS absorber toward Fairall9 is not the only example of a circumgalactic gas structure in the nearby Universe that exhibits a high overall metallicity together with a low N/ α abundance ratio. For example, Jenkins et al. (2005) have measured $[\alpha/H] = -0.19$ and $[N/\alpha] < 0.59$ in an intervening Lyman-limit system (LLS) at $z = 0.081$ toward the quasar PHL1811. This

absorption system appears to be associated with two nearby spiral galaxies at impact parameters $\rho = 34 h_{70}^{-1}$ and $\rho = 87 h_{70}^{-1}$ kpc. It possibly represents gaseous material that has been ejected or stripped from these galaxies (or other galaxies nearby) and thus possibly has an origin that is very similar to that of the Magellanic Stream. Note that low $[N/\alpha]$ ratios are also found in low-metallicity HVCs, such as Complex C (e.g., Richter et al. 2001b). Clearly, a systematic study of N/α ratios in circumgalactic metal absorbers at low z with HST/COS would be an important project to investigate, whether similar abundance patterns are typical for circumgalactic gaseous structures in the local Universe.

5.2. Physical conditions and small-scale structure in the gas

In addition to the very interesting chemical properties, the combined UVES/COS/FUSE/GASS data set provides a deep insight into the physical conditions in the Stream toward Fairall 9. The data show a complex multiphase gas structure that possibly spans a large range in gas densities and ionization conditions.

The presence of H_2 absorption toward Fairall 9 and NGC 3783 (Sembach et al. 2001) indicate that the MS and its Leading Arm hosts a widespread (because of the large absorption cross section/detection rate) cold neutral gas phase, possibly structured in a large amount of small, dense clumps or filaments. Assuming a total area of the MS and its Leading Arm of ~ 1500 deg², and a typical size for these clumps of a few pc, there could be millions of these dense structures in the neutral gas body of the Stream, if the two H_2 detections toward Fairall 9 and NGC 3783 reflects the true absorption-cross section of this gas phase throughout the Stream’s neutral gas body.

The fact that the MS can maintain significant amounts of H_2 at moderate gas densities ($n_H = 1 - 10$ cm⁻³) probably is a result of the relatively low intensity of the dissociating UV radiation at the Stream’s location due to the absence of local UV sources (see also Fox et al. 2005). Note that the low excitation temperature of H_2 for $J \geq 2$ (see also Sembach et al. 2001) provides independent evidence for a relatively weak ambient UV radiation field, and thus supports this scenario. In the MS, the low dissociation rate is expected to overcompensate the low formation rate (because of the lower metallicity and dust content compared to the Milky Way ISM), thus resulting in molecular gas fractions that are relatively high for the given total gas column. With a grain-formation rate of $R_{MS} = 3 \times 10^{-18}$ cm³ s⁻¹ and a density of $n_H = 5$ cm⁻³ (see Sect. 4.4), the H_2 formation time is long, $t_{form} = (R_{MS} n_H)^{-1} \approx 2$ Gyr, which matches the tidal age of the Stream. Therefore, it seems most likely, that the H_2 (or at least some fraction of it) has already formed in the parent galaxy and then survived the subsequent tidal stripping. A similar conclusion was drawn from the H_2 observation in the Leading Arm of the MS toward NGC 3783 (Sembach et al. 2001). Elaborating our idea, that the gas was locally enriched by massive stars shortly before the Stream was separated from its parent galaxy, one could imagine the presence of relatively dense, compressed shells and fragments that formed the seed structures for the formation of H_2 that

then were carried along with the Stream’s gaseous body.

Independent constraints on the physical conditions in the gas come from the observed Na I/Ca II ratio. We measure a Na I/Ca II column-density ratio of 0.23 in component 4, where the H_2 is expected to reside. In the Milky Way ISM, such a ratio is typical for the warm neutral medium (WNM), where $n_H \leq 10$ cm⁻³ and $T = 10^2 - 10^4$ K, typically, and Ca II serves a trace species (e.g., Crawford 1992). In such gas, and without dust depletion of Ca and Na, a nearly constant Na I/Ca II of 0.025 ratio would be expected from detailed ionization models of these ions (see Crawford 1992; Welty et al. 1996; Richter et al. 2011). If these numbers would apply also to the conditions in the MS, they would indicate relative dust depletions of Ca and Na of $\log \delta(Ca) - \log \delta(Na) \approx 0.9$, or $\log \delta(Ca) \approx 0.9$ for $\log \delta(Na) = 0$. These values are in very good agreement with the dust-depletion estimates from our Cloudy model discussed in Sect. 4.3.

5.3. Relevance to intervening QSO absorbers

The Magellanic Stream represents a prime example for a high-column density circumgalactic tidal gas stream in the local Universe. If the Stream would be seen as QSO absorber, it would be classified as LLS, sub-DLA (sub-Damped Lyman α absorber), or DLA, depending on the position of the sightline passing through the gas. The H I column-density map presented in Fig. 1 provides an estimate on the covering fractions of these column-density regime in the MS in the direction of Fairall 9. The results from our multi-sightline campaign to study the properties of the MS therefore are also of relevance for the interpretation of intervening absorption-line systems at low redshift.

5.3.1. Inhomogeneity of absorbers

The first important result from our study that is of relevance for QSO absorption-line studies is, that the physical conditions *and* the chemical composition appear to vary *substantially* within the Stream.

If such inhomogeneities were typical for tidal gas filaments around galaxies, then the interpretation of absorption spectra from circumgalactic gas structures around more distant galaxies (in terms of metallicity, molecular content, physical conditions, gas mass, origin, etc.) would be afflicted with large systematic uncertainties. In fact, most studies that aim at exploring the connection between galaxies and their surrounding circumgalactic gas are limited to single sightlines that pass the galaxy environment at a random impact parameter (e.g., Thom et al. 2011; Tumlinson et al. 2011; Ribaldo et al. 2011). Physical and chemical properties derived from single-sightline studies, however, may not be representative at all for the conditions in the general gaseous environment (in the same way as conditions in the MS along the Fairall 9 sightline are *not* representative for the Stream as a whole). The inhomogeneous chemical composition of the Stream implies that the metals possibly are not well mixed in the gas. Former abundance variations in the parent galaxy due to local star-formation events thus may have been frozen into the Stream’s spatial metal distribution. This would be not too surprising, however, since the main process that stir up and mix the interstellar gas within galaxies (i.e., supernova explosions, stellar

winds) cannot take place in the Stream simply due to the lack of stars.

5.3.2. Molecular absorption.

The two detections of H_2 absorption in the MS with *FUSE* toward Fairall9 and NGC 3783 (Sembach et al. 2001) indicate that tidal gas streams around galaxies may typically host a widespread, cold gas phase that has a substantial absorption cross section. This scenario is supported by the recent detection of H_2 absorption in another, more distant circumgalactic tidal gas stream beyond the Local Group (Crighton et al. 2013). These findings also remind us that the presence of H_2 in an intervening absorber with a complex velocity structure and a high neutral gas column density does not *necessarily* mean that one traces a gaseous disk of a galaxy.

The relatively large absorption cross section of H_2 in neutral gas structures *around* galaxies, as found in the Milky Way halo (see Richter 2006 and references therein), can be explained by the circumstance, that the physical conditions in these star-less gas clouds favour the formation of diffuse molecular structures even at relatively moderate gas densities. In particular, the relative low intensity of the local UV radiation field, the efficient process of H_2 line self-shielding, and the lack of the destructive influence of massive stars probably represent important aspects in this context.

H_2 observations in circumgalactic gas clouds not only provide important information on the physical and chemical conditions in these structures, they also are of high relevance for our understanding of the physics of molecular hydrogen in diffuse gas under conditions that are very different from that in the local ISM (see also Sembach et al. 2001). The transition from neutral to molecular gas, in particular, represents one of the most important processes that govern the evolutionary state of galaxies at low and high redshift. Detailed measurements of H_2 fractions and dust-depletion patterns in (star-less) tidal streams at low redshift can help to constrain the critical formation rate of molecular hydrogen in low-metallicity environments and thus could be of great importance to better understand the distribution and cross section of molecular gas in and around high-redshift galaxies.

6. SUMMARY

In this second paper of our ongoing series to study the Magellanic Stream in absorption we have analyzed newly obtained optical and UV absorption-line data from HST/COS and VLT/UVES together with archival *FUSE* and H I 21cm emission data from GASS to study the chemical composition and the physical conditions in the Magellanic Stream in the direction of the quasar Fairall9. Our main results are the following:

1. Metal absorption in the Magellanic Stream (MS) is detected in seven individual absorption components centered at $v_{\text{LSR}} = +143, +163, +182, +194, +204$ and $+218 \text{ km s}^{-1}$, indicating a complex internal velocity structure of the MS in this direction. Detected ions, atoms and molecules in the Stream include C IV , Si IV , Si III , C II , C II^* , Al II , Si II , S II , Ca II , Ti II , Fe II , Ti II , O I , N I , Na I , and H_2 .

2. From the unsaturated S II absorption and a Cloudy

photoionization model we measure an α abundance in the Stream of $[\text{S/H}] = -0.30 \pm 0.04$ (0.50 solar), which is substantially higher than that found in the Stream along the lines of sight toward NGC 7469, RBS 144, and NGC 7714 (Fox et al. 2010, 2013). Unfortunately, the unresolved, saturated $\text{O I } \lambda 1302.2$ line cannot be used to independently constrain the α abundance in the MS toward Fairall9. Contrary to sulfur, we measure a very low nitrogen abundance in the gas of $[\text{N/H}] = -1.15 \pm 0.06$. The resulting $[\text{N/S}]$ ratio is -0.85 dex, which is very low compared to other environments with similarly high α abundances. The low $[\text{N/S}]$ and high $[\text{S/H}]$ ratios observed in the MS toward Fairall9 suggest an abundance pattern that is dominated by the α enrichment from massive stars and Type II supernovae, while only very little primary nitrogen was deposited into the gas when the Stream was separated from the Magellanic Clouds.

3. The detection of very narrow Na I and H_2 absorption (with $b \approx 2 \text{ km s}^{-1}$) in the component at $v_{\text{LSR}} = +188 \text{ km s}^{-1}$ indicates the presence of a compact (pc-scale), cold gas structure in the MS along this sightline. From the analysis of the newly reduced archival *FUSE* data of Fairall9 we measure a total molecular hydrogen column density of $\log N(\text{H}_2) = 17.93^{+0.19}_{-0.16}$, improving previous results from Richter et al. (2001a). From the analysis of the H_2 rotational excitation we obtain a kinetic temperature in the cold neutral gas phase of $T \approx 93 \text{ K}$. For the gas density we derive $n \approx 4 - 8 \text{ cm}^{-3}$, assuming that the H_2 gas is in a formation-dissociation equilibrium. The resulting estimate for the thermal gas pressure is $P/k \approx 750 \text{ cm}^{-3} \text{ K}$, thus in good agreement with values derived from previous studies of the Stream. The detection of H_2 absorbing structures in the MS, whose linear and angular sizes must be very small ($\sim 1 \text{ pc}$ and $\sim 3 \text{ arcseconds}$), indicates that the neutral gas body of the Stream is highly clumped and structured.

4. The relatively low column densities of Fe II , Ti II , Ni II , Al II , Ca II indicate that the gas phase abundances of these elements are affected by dust depletion. We combine our column-density measurements for these ions with a Cloudy photoionization model and derive dust depletion values relative to sulfur of $\log \delta(\text{Si}) = -0.27$, $\log \delta(\text{Fe}) = -0.56$, $\log \delta(\text{Ni}) \leq -0.51$, $\log \delta(\text{Ti}) = -1.32$, $\log \delta(\text{Al}) = -0.62$, and $\log \delta(\text{Ca}) = 0$ to -1.58 . These depletion values are similar to those found in warm, diffuse clouds in the lower Milky Way halo (Savage & Sembach 1996).

5. Our study indicates that the enrichment history of the Magellanic Stream as well as the physical conditions in the Stream are more complex than previously known. The abundances and gas-to-dust ratios measured in the Stream along the Fairall9 sightline are substantially higher than what is found along other MS sightlines. The low N abundance favours an SMC origin of the MS gas toward Fairall9, although an LMC origin cannot be ruled out. The high sulfur abundance in the gas possibly indicates a massive α enrichment from massive stars in a region of enhanced star-formation $\sim 1 - 2 \text{ Gyr}$ ago and then was stripped from the Magellanic Clouds

and incorporated into the MS, before the gas could be enriched in nitrogen from intermediate stars.

We gratefully thank Gurtina Besla and Mary Putman for helpful discussions. Support for this research was provided by NASA through grant HST-GO-12604 from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555.

REFERENCES

- Abgrall, H., & Roueff, E. 1989, *A&A*, 79, 313
- Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Asplund, M., Grevesse, N., Jacques Sauval, A., & Scott, P. 2009, *ARA&A*, 47, 481
- Ben Bekhti, N., Richter, P., Westmeier, T., & Murphy, M.T. 2008, *A&A*, 487, 583
- Ben Bekhti, N., Winkel, B., Richter, P., Kerp, J., Klein, U., Murphy, M. T. 2011, *A&A*, 542, 110
- Besla, G., Kallivayalil, N., Hernquist, L., et al. 2007, *ApJ*, 668, 949
- Besla, G., Kallivayalil, N., Hernquist, L., et al. 2010, *ApJ*, 721, L97
- Besla, G., Kallivayalil, N., Hernquist, L., van der Marel, R. P., Cox, T. J., & Keres, D. 2012, *MNRAS*, 421, 2109
- Bland-Hawthorn, J., & Maloney, P. R. 1999, *ApJ*, 510, L33
- Bland-Hawthorn, J., Sutherland, R., Agertz, O., & Moore, B. 2007, *ApJ*, 670, L109
- Bland-Hawthorn, J., & Maloney, P. R. 2002, in *ASP Conf. Ser.* 254, *Extragalactic Gas at Low Redshift*, ed. J. S. Mulchaey & J. T. Stocke (San Francisco: ASP), 267
- Brüns, C., Kerp, J., Staveley-Smith, L., et al. 2005, *A&A*, 432, 45
- Crawford, I.A. 1992, *MNRAS*, 259, 47
- Crighton, N. H. M., Bechtold, J., Carswell, R. F. et al. 2013, *MNRAS*, in press (astro-ph/1210.0905)
- Connors, T. W., Kawata, D., Maddison, S. T., & Gibson, B. K. 2004, *PASA*, 21, 222
- Connors, T. W., Kawata, D., & Gibson, B. K. 2006, *MNRAS*, 371, 108
- Dalgarno, A., & McCray, R. A. 1972, *ARA&A*, 10, 375
- de Boer, K. S., Richter, P., Bomans, D. J., Heithausen, A., & Koornneef, J. A&A, 338, L5
- Dekker, H., D'Odorico, S., Kaufer, A., Delabre, B., & Kotzlowski, H. 2000, *SPIE*, 4008, 534
- Diaz, J. D., & Bekki, K. 2011, *ApJ*, 413, 2015
- Draine, B., & Bertoldi, F. 1996, *ApJ*, 468, 269
- Dufour, R.J. 1975, *ApJ*, 195, 315
- Ferland, G. J., Korista, K. T., Verner, D. A., Ferguson, J. W., Kingdon, J. B., & Verner, E. M. 1998, *PASP*, 110, 761
- Fontana, A., & Ballester, P. 1995, *ESO Messenger*, 80, 37
- Fox, A. J., Wakker, B. P., Savage, B. D., Tripp, T. M., Sembach, K. R., & Bland-Hawthorn, J. 2005, *ApJ*, 630, 332
- Fox, A.J., Wakker, B.P., Smoker, J.V., Richter, P., Savage, B.D., & Sembach, K.R. 2010, *ApJ*, 718, 1046
- Fox, A.J., Richter, P., Wakker, B.P., Lehner, N., Howk, J.C., Ben Bekhti, N., Bland-Hawthorn, J., & Lucas, S. 2013, *ApJ*, in press
- Gardiner, L.T. & Noguchi, M. 1996, *MNRAS*, 278, 191
- Gibson, B. K., Giroux, M. L., Penton, S. V., Putman, M. E., Stocke, J. T., & Shull, J. M. 2000, *AJ*, 120, 1830
- Green, J. C., Froning, C. S., Osterman, S., et al. 2012, *ApJ*, 744, 60
- Harris, J., Zaritsky, D. 2004, *AJ*, 127, 1531
- Harris, J., Zaritsky, D. 2009, *AJ*, 138, 1243
- Henry, R. B. C., Edmunds, M. G., Köppen, J. 2000, *ApJ*, 541, 660
- Henry, R. B. C. & Prochaska, J. X. 2007, *PASP*, 119, 962
- Hughes, J. P., Hayashi, I., & Koyama, K. 1998, *ApJ*, 505, 732
- Irwin, M. J., Kunkel, W. E., Demers, S. 1985, *Nature*, 318, 160
- Jenkins, E. B., Bowen, D. V., Tripp, T. M., & Sembach, K. R. 2005, *ApJ*, 623, 767
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., Arnal, E. M., Bajaja, E., Morras, R., & Pöppel, W. G. L. 2005, *A&A*, 440, 775
- Kalberla, P. M. W., McClure-Griffiths, N. M., Pisano, D. J., et al. 2010, *A&A*, 521, 17
- Kallivayalil, N., van der Marel, R. P., Alcock, C., Axelrod, T., Cook, K. H., Drake, A. J., & Geha, M. 2006a, *ApJ*, 638, 772
- Kallivayalil, N., van der Marel, R. P., & Alcock, C. 2006b, *ApJ*, 652, 1213
- Keenan, F. P., Lennon, D. J., Johnson, C. T., & Kingston, A. E. 1986, *MNRAS*, 220, 571
- Kobayashi, C., Umeda, H., Nomoto, K., Tominaga, N., Ohkubo, T. 2006, *ApJ*, 653, 1145
- Koerwer, J.F. 2009, *AJ*, 138, 1
- Kriss, G. A. 2011, *COS Instrument Science Report*, 1
- Lehner, N. 2002, *ApJ*, 578, 126
- Lehner, N., Wakker, B.P., & Savage, B.D. 2004, *ApJ*, 615, 767
- Lehner, N., Howk, J.C., Keenan, F.P., & Smoker, J.V. 2008, *ApJ*, 678, 219
- Lu, L., Savage, B. D., Sembach, K. R. 1994, *ApJ*, 437, L119
- Mastropietro, C., Moore, B., Mayer, L., Wadsley, J., & Stadel, J. 2005, *MNRAS*, 363, 509
- Mathewson, D. S., Ford, V. L., Schwarz, M. P., & Murray, J. D. 1979, *IAUS*, 84, 547
- Melnick, J. 1985, *A&A*, 153, 235
- Morton, D. C., Dinerstein, H. L. 1976, *ApJ*, 204, 1
- Morton, D. C. 2003, *ApJS*, 149, 205
- McClure-Griffiths, N. M., Pisano, D. J., Calabretta, M. R., et al. 2009, *ApJS*, 181, 398
- Nidever, D. L., Majewski, S. R., & Burton, W. B. 2008, *ApJ*, 679, 432
- Nidever, D. L., Majewski, S. R., Burton, W. B., & Nigra, L. 2010, *ApJ*, 723, 1618
- Nigra, L., Stanimirovic, S., Gallagher, J. S., Wood, K., Nidever, D., & Majewski S. 2013, *ApJ*, 760, 48
- Pagel, B. E. J., & Tautvaisiene, G. 1998, *MNRAS*, 299, 535
- Pettini, M., Ellison, S. L., Bergeron, J., & Petitjean, P. 2002, *A&A*, 391, 21
- Pettini, M., Zych, B. J., Steidel, C. C., & Chaffee, F. H. 2008, *MNRAS*, 385, 2011
- Putman, M. E., Staveley-Smith, L., Freeman, K. C., Gibson, B. K., & Barnes, D. G. 2003, *ApJ*, 586, 170
- Ribaudo, J., Lehner, N., Howk, J. C., Werk, J., Tripp, T. M., Prochaska, J. X., Meiring, J. D. & Tumlinson J. 2011, *ApJ*, 743, 207
- Richter, P., Widmann, H., de Boer, K. S. et al. 1998, *A&A*, 338, L9
- Richter, P. 2000, *A&A*, 359, 1111
- Richter, P., Sembach, K. R., Wakker, B. P., Savage, B. D., 2001a, *ApJ*, 562, L181
- Richter, P., Savage, B.D., Wakker, B.P., Sembach, K.R., Kalberla, P.M.W. 2001b, *ApJ*, 549, 281
- Richter P., Wakker B.P., Savage B.D., & Sembach K.R. 2003, *ApJ*, 586, 230
- Richter, P., Westmeier, T., & Brüns 2005, *A&A*, 442, L49
- Richter, P. 2006, *Rev. Mod. Astron.*, 19, 31
- Richter, P., Charlton, J.C., Fangano, A.P.M., Ben Bekhti, N., & Masiero, J.R. 2009, *ApJ*, 695, 1631
- Richter, P., Krause, F., Fechner, C., Charlton, J.C., & Murphy, M.T. 2011, *A&A*, 528, A12
- Russell, S. C., & Dopita, M. A. 1990, *ApJ*, 74, 93
- Russell, S. C., & Dopita, M. A. 1992, *ApJ*, 384, 508
- Savage, B. D., Drake, J. F., Budich, W., & Bohlin, R. C. 1977, *ApJ*, 216, 291
- Savage, B. D., & Sembach, K. R. 1991, *ApJ*, 379, 245
- Savage, B.D. & Sembach, K.R. 1996, *ARA&A*, 34, 279
- Sembach, K.R., Howk, J.C., Savage, B.D., Shull, J.M. 2001, *AJ*, 121, 992
- Songaila, A. 1981, *ApJ*, 243, L19
- Spitzer, L. 1978, *Physical processes in the interstellar medium*, (New York Wiley-Interscience)
- Thom, C., Werk, J. K., Tumlinson, J., Prochaska, J. X., Meiring, J. D., Tripp, T. M., & Sembach, K. R. 2011, *ApJ*, 736, 1
- Tumlinson, J., Shull, J. M., Rachford, B. L., et al. 2002, *ApJ*, 566, 857
- Tumlinson, J., Werk, J. K., Thom, C., et al. 2011, *ApJ*, 733, 111
- Wakker, B. P., Howk, J. C., Savage, B. D., et al. 1999, *Nature*, 402, 388
- Wakker, B. P., Oosterloo, T. A., Putman, M. E. 2002, *AJ*, 123, 1953
- Wakker, B.P. 2006, *ApJS*, 163, 282
- Wakker, B. P., York, D. G., Howk, J. C., et al. 2007, *ApJ*, 670, L113
- Wakker, B. P., York, D. G., Wilhelm, R., Barentine, J. C., Richter, P., Beers, T. C., Ivezić, Z., & Howk, J.C. 2008, *ApJ*, 672, 298
- Wakker, B. P., Lockman, F. J., & Brown, J. M. 2011, *ApJ*, 728, 159
- Wannier, P., & Wrixon, G. T. 1972, *ApJ*, 173, L119
- Weiner, B. J., & Williams, T. B. 1996, *AJ*, 111, 1156
- Weisz, D. R., Doplin, A. E., Skillmann, E. D., et al. 2013, *MNRAS*, in press (aXiv:1301.7422)
- Welty, D. E., Morton, D. C., & Hobbs, L.M. 1996, *ApJS*, 106, 533
- Welty, D. E., Lauroesch, J. T., Blades, J. C., Hobbs, L.M., & York, D. G. 1997, *ApJ*, 489, 672
- Welty, D. E., Federman, S. R., Gredel, R., Thorburn, J. A., & Lambert, D. L. 2006, *ApJS*, 165, 138
- Wolfire, M. G., McKee, C. F., Hollenbach, D., & Tielens, A. G. G. M. 1995, *ApJ*, 453, 673
- York, D. G., Songaila, A., Blades, J. C., Cowie, L. L., Morton, D. C., & Wu, C.-C. 1982, *ApJ*, 255, 467
- Zech, W. F., Lehner, N., Howk, J. C., Dixon, W. V. D., & Brown, T. M. 2008, *ApJ*, 679, 460




FIG. 10.— H I Ly α absorption in the COS spectrum of Fairall 9 in the wavelength range between 1195 and 1235 Å. The red solid line indicates a two-component absorption model with a Milky Way H I column density of $\log N(\text{H I}) = 20.25$ and a MS H I column density of $\log N(\text{H I}) = 19.95$.

APPENDIX

H I LY α ABSORPTION

In Fig. 10 we show the normalized COS spectrum in the wavelength range between 1195 and 1235 Å. Strong H I Ly α absorption centered at 1216 Å from neutral gas in the Milky Way (MW) and in the Magellanic Stream with well-defined damping wings is visible. The 21cm data from various observations and instruments (Sect. 3.3; Gibson et al. 2000) indicate that the Milky Way disk gas dominates the neutral-gas column density along this sightline

The strength and shape of the Ly α absorption toward Fairall 9 supports this conclusion, but the relative contribution of MW and MS cannot be tightly constrained from a Voigt-profile fit of the Ly α absorption. The red solid line in Fig. 10 shows a two-component absorption model with a MW H I column density of $\log N(\text{H I}) = 20.25$ and a MS H I column density of $\log N(\text{H I}) = 19.95$. Therefore, the Ly α absorption profile is fully consistent with the value of $\log N(\text{H I}) = 19.95$ in MS, as indicated by the GASS 21cm data. However, the Ly α profile is also consistent with MS column densities in the range $\log N(\text{H I}) = 19.75 - 20.15$, if the MW contribution is modified accordingly. In conclusion, the Ly α absorption does not provide constraining limits on the H I column density in the Magellanic Stream toward Fairall 9 that could help to minimize the systematic error for the determination of $\log N(\text{H I})$, as discussed in Sect. 3.3.

ABSORPTION MODELING OF UV LINES

In Sect. 2.5 we have outlined the method that we have used to model the UV lines of the low ions in the COS spectrum of Fairall 9 to infer column densities of these species. We here present additional information on the modeling procedure to provide an insight into the allowed parameter range for (N_i, b_i) and N_{tot} for each ion.

In Fig. 11 we show velocity plots of the low ions from synthetic spectra that have been generated by us based on different input models, in which we have systematically varied the parameters N_i and b_i . The resulting model profiles are compared to the best-fitting model presented in Fig. 3 and Table 1.

In the left panel of Fig. 11 (a), the best-fitting model is indicated with the black solid line, while the lines in red, green, and blue show models, in which all column densities N_i in the individual subcomponents have been increased by 0.3, 0.5, and 1.0 dex, respectively. The velocity structure and the b values remain unchanged. In the middle panel of Fig. 11 (b), the column densities N_i in the individual subcomponents have been decreased by 0.3, 0.5, and 1.0 dex (red, green, blue), again without changes in v_i and b_i . These two figures indicate that the shapes of the strong and saturated lines from C II, O I and Si II are basically not affected by the change in column density. In contrast, the absorption depths in the weaker lines of N I, Al II, Si II, S II, and Fe II are *strongly* affected by the increase or decrease of the column densities with respect to the reference model (black), demonstrating that these lines represent sensitive diagnostics to constrain N_i and N_{tot} for these ions in the Stream.

In the right panel of Fig. 11 (c) we show a model in which we have increased the Doppler parameters b_i for the individual subcomponents by factors of 1.2 (red), 1.5 (green), and 2.0 (blue) with respect to the reference model (black), while the column densities and the overall velocity structure remain unchanged. Because of the relatively low spectral resolution of COS compared to the absolute values of b_i , the changes in b_i are irrelevant for the shape of the weak, unsaturated lines, while the strong, saturated lines mildly grow in strength in their absorption wings. Therefore, even if we would underestimate the values of b_i for the singly-ionized species Al II, Si II, S II, and Fe II when adopting the values from Ca II, as one could argue because these ions may trace a slightly different (more extended) gas phase, our column-density estimate would not be affected at all by this systematic error.

These figures, together with Fig. 2, underline, that the total gas column densities of N I, Al II, Si II, S II, and Fe II in the MS, as listed in Table 1, are well constrained by the COS data and our absorption model.




FIG. 11.— Velocity profiles of low ions, as derived from synthetic spectra from different input models for the gas absorption in the Magellanic Stream toward Fairall 9. Details are given in the text in the Appendix.